

# Perinatal Statistics Report

Health Research and Information Division
October 2009

2007



#### **ACKNOWLEDGEMENTS**

The National Perinatal Reporting System (NPRS) within the Health Research and Information Division (HRID) at the Economic and Social Research Institute (ESRI) is responsible for collecting, coding, inputting, and validating data on all births nationally on an annual basis. Data on 71,963 births were analysed for this report. For each of these records additional information, apart from that required to register each birth, had to be collected in each hospital and by each independent midwife.

The collection and analysis of national perinatal data is a large undertaking, and the compilation of an accurate and comprehensive database on a timely basis is greatly dependent on the commitment and support of those at the forefront of collecting the data. Principal among these are the staff of the maternity hospitals and units, as well as the many independent midwives involved in practice. We would like to express our sincere appreciation to all who contribute to the accurate recording and timely notification of statistical information for all births nationally.

In addition to financing the NPRS, the Department of Health and Children and the Health Service Executive continues to provide essential support for the operation and development of this system. We would like to acknowledge Hugh Magee and Alan Cahill, in particular, for the advice and assistance provided in the course of preparing this report.

The production of this report is very much a team effort within the Health Research and Information Division. We would like to acknowledge, specifically, Aoife Brick, Richard Layte, Aisling Mulligan, Deirdre Murphy, Sinead O'Hara, Jacqueline O'Reilly, and Eithne Sexton for reviewing earlier drafts of this report. We would also like to thank Brian McCarthy and Shane McDermott for their work on the development of the NPRS software system for data entry and validation. Preparation of the data and production of this report was undertaken by Sheelagh Bonham, assisted by Barbara Clyne, Catherine Glennon, Anna Lachacz, and Izabella Nowacka. We wish to express our sincere thanks to all our colleagues for their hard work and commitment.

### **TABLE OF CONTENTS**

LIST OF TABLES	
LIST OF FIGURES	
KEY FINDINGS	
SECTION ONE	
INTRODUCTION	
Introduction	
Purpose of this Report	
Structure of this Report	
Conventions used in this Report	
National Minimum Data Set	
Data Collection and Quality	
SECTION TWO	
TIME TRENDS AND INTERNATIONAL COMPARISONS	
Introduction	
Trends in Selected Statistics, 1999-2007	
Perinatal Mortality	
Birth Rates	
Fertility	
Singleton and Multiple Births	
SECTION TUDES	,
SECTION THREE	1
GENERAL CHARACTERISTICS	
Introduction	
Maternal Age	
Mother's Marital Status	
Occupation	
Mother's Occupation	
Father's Occupation	
Nationality	
Mother's Nationality	
Father's Nationality	
Maternal Parity	
Number of Previous Stillbirths	
Interval Since Last Birth	
Birthweight	
Gestational Age	
Month of Birth	

SECTION FOUR	
PERINATAL CARE	
Introduction	
Type of Antenatal Care	
Duration of Pregnancy at First Antenatal Visit	
Mother's Antenatal Length of Stay	
Mother's Postnatal Length of Stay	
Advance Hospital Bookings	
Rubella: Immune Status of Mother	
Method of Delivery	
Size of Maternity Unit	
Inter-Hospital Transfer of Infants	
Infant's Length of Stay	
Infant's Type of Feeding	
SECTION FIVE	
PERINATAL OUTCOMES	
Introduction	
Mortality by Birthweight Category	
Cause of Death by Birthweight	
Mortality by Gestational Age	
Cause of Death	
Age at Death	
Post-Mortem Examination Uptake	
APPENDICES	
Appendix A: Definitions	
Appendix B: Copy of Part 3 of the Birth Notification Form	
Appendix C: Classification of Occupation	
Appendix D: Cause of Death Classification	
Appendix E: Classification of County of Residence	
Appendix F: Classification of Nationality	
Appendix G: Domiciliary Births, 1999-2007	
Appendix H: Source Data for Figures	
Appendix 1: Table Reference Guide to Previous Perinatal Statistics Report 2006	

#### **LIST OF TABLES**

2.1	Trends in Selected Statistics, 1999-2007	9
2.2	Live Births, Stillbirths, Early Neonatal Deaths, and Perinatal Deaths,	11
	Ireland, 1999-2007	
2.3	Singleton and Multiple Births, 1999-2007	17
2.4	Live Births, Stillbirths and Early Neonatal Deaths, Numbers and Rates	17
	for Singleton and Multiple Births, 1999-2007	
Tables for Gene	eral Characteristics for the year 2007:	
3.1a	Age of Mother, Singleton Births	21
3.1b	Age of Mother, Multiple Births	21
3.2a	Mother's Marital Status, Singleton Births	23
3.2b	Mother's Marital Status, Multiple Births	23
3.3a	Mother's Occupation, Singleton Births	25
3.3b	Mother's Occupation, Multiple Births	26
3.4a	Father's Occupation, Singleton Births	28
3.4b	Father's Occupation, Multiple Births	29
3.5a	Mother's Nationality, Singleton Births	32
3.5b	Mother's Nationality, Multiple Births	33
3.6a	Father's Nationality, Singleton Births	36
3.6b	Father's Nationality, Multiple Births	37
3.7a	Maternal Parity, Singleton Births	38
3.7b	Maternal Parity, Multiple Births	39
3.8a	Previous Stillbirths, Singleton Births	41
3.8b	Previous Stillbirths, Multiple Births	41
3.9a	Interval in Years Since Last Birth, Singleton Births	42
3.9b	Interval in Years Since Last Birth, Multiple Births	42
3.10a	Birthweight, Singleton Births	43
3.10b	Birthweight, Multiple Births	44
3.11a	Gestational Age at Delivery, Singleton Births	46
3.11b	Gestational Age at Delivery, Multiple Births	46
3.12a	Month of Birth, Singleton Births	47
3.12b	Month of Birth, Multiple Births	48
	natal Care for the year 2007:	
4.1a	Type of Antenatal Care, Singleton Births	51
4.1b	Type of Antenatal Care, Multiple Births	51
4.2a	Time of First Visit to Doctor or Hospital during Pregnancy,	52
	Singleton Births	
4.2b	Time of First Visit to Doctor or Hospital during Pregnancy,	52
	Multiple Births	
4.3a	Mother's Antenatal Length of Stay, Singleton Births	53
4.3b	Mother's Antenatal Length of Stay, Multiple Births	53
4.4a	Mother's Postnatal Length of Stay, Singleton Births	54
4.4b	Mother's Postnatal Length of Stay, Multiple Births	55

4.5a	Advance Hospital Bookings, Singleton Births	5/
4.5b	Advance Hospital Bookings, Multiple Births	57
4.6a	Rubella: Immune Status of Mother, Numbers and Percentages,	58
	Singleton Births	
4.6b	Rubella: Immune Status of Mother, Numbers and Percentages,	58
	Multiple Births	
4.7a	Method of Delivery, Singleton Births	59
4.7b	Method of Delivery, Multiple Births	59
4.8	Method of Delivery by Mother's Age, Total Births	60
4.9	Size of Maternity Unit (number of live births and stillbirths annually)	61
4.10a	Inter-Hospital Transfer of Infants, Singleton Births	62
4.10b	Inter-Hospital Transfer of Infants, Multiple Births	62
4.11a	Percentage Distribution of Live Births by Infant's Length of Stay in	63
	Hospital, Singleton Births	
4.11b	Percentage Distribution of Live Births by Infant's Length of Stay in	63
	Hospital, Multiple Births	
4.12a	Age of Mother by Infant's Type of Feeding, Numbers and Percentages,	64
	Singleton Births	
4.12b	Age of Mother by Infant's Type of Feeding, Numbers and Percentages,	65
	Multiple Births	
4.13	Mother's Occupation by Infant's Type of Feeding, Total Live Births	67
Tables for I	Perinatal Outcomes for the year 2007:	
	Cimatal Catalonics for the year 2007.	
<b>5</b> 4 .	Line Birthe Antonia and Laborator of Cilliante Ford Norward	74
5.1a	Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal	71
Г 1 Ь	Deaths, and Mortality Rates by Birthweight, Singleton Births	72
5.1b	Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal	72
Г Эс	Deaths, and Mortality Rates by Birthweight, Multiple Births	75
5.2a	Perinatal Mortality: Numbers and Rates, by Birthweight and	75
F 21-	Cause of Death, Singleton Births	77
5.2b	Perinatal Mortality: Numbers and Rates, by Birthweight and	//
F 2-	Cause of Death, Multiple Births	70
5.3a	Live Births, Antepartum, and Intrapartum Stillbirths, Early Neonatal	79
г 2h	Deaths and Mortality Rates by Gestational Age, Singleton Births	00
5.3b	Live Births, Antepartum, and Intrapartum Stillbirths, Early Neonatal	80
<b>5</b> 4-	Deaths and Mortality Rates by Gestational Age, Multiple Births	0.2
5.4a	Stillbirths, Early Neonatal Deaths and Mortality Rates by Cause of	83
E 41.	Death by ICD-10 Chapter and Selected Sub-Categories, Singleton Births	0.5
5.4b	Stillbirths, Early Neonatal Deaths and Mortality Rates by Cause of	85
	Death by ICD-10 Chapter, Multiple Births	
	Age at Death, Early Neonatal Deaths, Singleton Births	89
5.5a		00
5.5b	Age at Death, Early Neonatal Deaths, Multiple Births	89
	Age at Death, Early Neonatal Deaths, Multiple Births Stillbirths and Early Neonatal Deaths undergoing	89 90
5.5b 5.6a	Age at Death, Early Neonatal Deaths, Multiple Births Stillbirths and Early Neonatal Deaths undergoing Post-Mortem Examinations, Singleton Births	90
5.5b	Age at Death, Early Neonatal Deaths, Multiple Births Stillbirths and Early Neonatal Deaths undergoing	

D1 D2	Cause of Death Coding Rules ICD-10, Cause of Death Codes	101 102
Tables for I	Domiciliary Births:	
G1	Trends in Selected Statistics, Domiciliary Births, 1999-2007	110
G2	Age of Mother by Maternal Parity, Domiciliary Births	111
G3	Age of Mother by Gestational Age at Delivery, Domiciliary Births	111
G4	Birthweight by Gestational Age at Delivery, Domiciliary Births	112
G5	Mother's Marital Status by Age of Mother, Domiciliary Births	112
G6	Mother's Occupation by Birthweight, Domiciliary Births	113
G7	Father's Occupation by Infant's Type of Feeding, Live Domiciliary Birth	
G8	Mother's Occupation by Infant's Type of Feeding, Live Domiciliary Birtl	
Tables for S	Source Data for Figures:	
H1	Age of Mother, Live Births, Stillbirths, Early Neonatal Deaths,	116
	and Mortality Rates, Total Births	
H2	Percentage of births by Mother's Marital Status, Total Births, 2003-200	
H3	Mother's County of Residence by Mother's Marital Status, Total Births	
H4	Mother's Occupation, Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, Total Births	119
H5	Father's Occupation, Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, Total Births	120
Н6	Percentage of Births by Mother's Nationality, Total Births, 2004-2007	121
H7	Mother's Average Age by Mother's Nationality, Total Births	121
H8	Percentage of Births to Primiparous Women by Maternal Age,	121
110	Total Births, 2003 and 2007	121
H9	Percentage of Births to Multiparous Women by Maternal Age,	122
	Total Births, 2003 and 2007	
H10	Age of Mother by Maternal Parity, Total Births	122
H11	Birthweight Groups, Numbers and Percentages, Total Live Births	122
H12	Birthweight Groups by Sex, Total Live Births	123
H13	Gestational Age at Delivery, Live Births, Stillbirths, Early Neonatal	123
	Deaths, and Mortality Rates, Total Births	40.
H14	Method of Delivery by Mother's Postnatal Length of Stay,	124
	Singleton Births	405
H15	Method of Delivery by Mother's Postnatal Length of Stay,	125
114.6	Multiple Births	405
H16	Age of Mother by Infant's Type of Feeding, Total Live Births	125
H17	Mother's County of Residence by Infant's Type of Feeding,	126
	Total Live Births	400
H18	Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal	128
114.0	Deaths, and Mortality Rates by Birthweight, Total Births	400
H19	Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal	128
	Deaths, and Mortality Rates by Gestational Age, Total Births	

#### **LIST OF FIGURES**

1.1	National Perinatal Reporting System, Data Collection	6
2.1	Stillbirths, Early Neonatal and Perinatal Mortality Rates, Ireland, 1999-2007	12
2.2	Perinatal Mortality Rates for Selected EU Countries, 2007	13
2.3	Trends in Birth Rates for Selected EU Countries, 1999-2007	14
2.4	Birth Rates per 1,000 Population for EU Countries, 2007	14
2.5	Age-Specific Fertility Rates (ASFRs) and Total Period Fertility	15
	Rates (TPFRs), Ireland, 1999-2007	
2.6	Total Period Fertility Rates for Selected EU Countries, 2007	16
3.1	Age-specific Stillbirth, Early Neonatal and Perinatal Mortality Rates, Total Births, 2007	22
3.2	Percentage of Births by Mother's Marital Status, Total Births, 2003-2007	7 24
3.3	Percentage of Births to Single Mothers by Mother's County of Residence, Total Births, 2007	24
3.4	Perinatal Mortality Rates by Mother's Occupation, Total Births, 2007	27
3.5	Perinatal Mortality Rates by Father's Occupation, Total Births, 2007	30
3.6	Percentage of Births by Nationality for Non-Irish Mothers, Total Births, 2004-2007	34
3.7	Average Age of Mother by Mother's Nationality, Total Births, 2007	34
3.8	Total Births to Primiparous Women in each Maternal Age Group, 2003 and 2007	39
3.9	Total Births to Multiparous Women in each Maternal Age Group, 2003 and 2007	40
3.10	Parity by Age of Mother, Total Births, 2007	40
3.11	Birthweights: Cumulative Percentages of Total Live Births, 2007	44
3.12	Percentage Births by Birthweight Groups, Total Live Births, 2007	45
3.13	Percentage Distribution for Males and Females across Birthweight Groups, Total Births, 2007	45
3.14	Perinatal Mortality Rates by Gestational Age at Delivery, Total Births, 2007	47
3.15	Percentage of Births by Month of Birth, Total Births, 2007	48
4.1	Mother's Postnatal Length of Stay by Method of Delivery,	56
	Singleton Births, 2007	
4.2	Mother's Postnatal Length of Stay by Method of Delivery,	56
	Multiple Births, 2007	
4.3	Percentage of Births by Method of Delivery for Live Singleton and Live Multiple Births, 2007	60
4.4	Percentage of Births by Size of Maternity Unit, Total Live Births, 2007	61
4.5	Percentage of Births by Infant's Length of Stay for Live Singleton and Live Multiple Births, 2007	63

4.6 Percentage of Breastfed Infants by Age of Mother, Total Live Births, 2007					
4.7	Percentage Distribution of Breastfed Infants by Mother's County of Residence, Total Live Births, 2007	66			
4.8	Percentage Distribution of Infant's Feeding by Mother's Occupation, Total Live Births, 2007	68			
5.1	Deaths by Birthweight Groups, Singleton Births, 2007	72			
5.2	Deaths by Birthweight Groups, Multiple Births, 2007	73			
5.3	Perinatal Mortality Rates by Birthweight Groups, Total Births, 2007	73			
5.4	Total Deaths < 2,500 grams by Cause of Death, 2007	78			
5.5	Total Deaths ≥ 2,500 grams by Cause of Death, 2007	78			
5.6	Deaths by Gestational Age Groups, Singleton Births, 2007	80			
5.7	Deaths by Gestational Age Groups, Multiple Births, 2007	81			
5.8	Perinatal Mortality Rates by Gestational Age at Delivery Total Births, 2007	81			
5.9	Stillbirths by Cause of Death, Singleton Births, 2007	86			
5.10	Stillbirths by Cause of Death, Multiple Births, 2007	86			
5.11	Early Neonatal Deaths by Cause of Death, Singleton Births, 2007	87			
5.12	Early Neonatal Deaths by Cause of Death, Multiple Births, 2007	87			
5.13	Perinatal Deaths by Cause of Death, Singleton Births, 2007	88			
5.14	Perinatal Deaths by Cause of Death, Multiple Births, 2007	88			

#### **KEY FINDINGS**

The Perinatal Statistics Report 2007 presents national statistics on perinatal events in Ireland. This report is based on data collected from the National Perinatal Reporting System (NPRS) and provides information on mothers giving birth and babies born in 2007.

#### **SUMMARY STATISTICS, 2007**

Live Births, Stillbirths, Early Neonatal Deaths, Perinatal Deaths, and Adjusted Perinatal Deaths

	Total Births	Live Bii	rths	Stillbirths		Early Neonatal Deaths		Perinatal Deaths		Adjusted Perinatal Deaths*	
	Number	Number	Rate	Number	Rate	Number	Rate	Number	Rate	Number	Rate
Singleton	69,692	69,318	-	374	5.4	118	1.7	492	7.1	354	5.1
Multiple	2,271	2,252	-	19	8.4	14	6.2	33	14.5	29	12.8
Total	71,963	71,570	-	393	5.5	132	1.8	525	7.3	383	5.4

<sup>\*</sup> In this and subsequent tables, 'Adjusted Perinatal Rate' excludes all deaths due to congenital anomalies.

#### **BABIES:**

#### Of the babies born in 2007

- 6.1% were preterm (less than 37 weeks gestation), the highest reported since 1999 (5.7%)
- 5.3% of live births were low birthweight (less than 2,500 grams)
- 45% of babies were breastfed at discharge from hospital, this compares to 41% in 2003 and 36% in 1999
- 26% of total live births were delivered by Caesarean section. This rate was 25% for singletons and 64% for multiples. The Caesarean section rate for total live births was 20% in 1999
- the average length of stay in hospital was 3.3 days for a singleton birth and 10.0 days for a multiple birth
- the perinatal mortality rate was 7.3 per 1,000 live births and stillbirths; 7.1 for singleton births and 14.5 for multiple births; for the years 1999-2007 the highest perinatal mortality rate was 8.9 in 2000 and the lowest recorded was 7.0 in 2006
- a perinatal mortality rate of 8.7 per 1,000 live births and stillbirths was recorded for babies to mothers aged 19 years or less and to those aged 35 years or more, which was higher than the perinatal mortality rates recorded for babies born to mothers in the other age group categories.

#### **MOTHERS:**

#### Of the women who gave birth in 2007

- average age has steadily increased from 30.1 years in 1999 to 31.1 years in 2007
- 27% were aged 35 years or older, up from 21% in 1999
- 14% of first births were to women aged 35 years or older, this has been increasing each year and was recorded at 9% for 1999
- 3.5% of total mothers giving birth were aged 19 years or less, compared to 6.2 % in 1999
- 42% gave birth for the first time, with an average age for first time mothers of 28.8 years
- 32% of births were to single mothers, with an average age for single mothers of 27.1 years
- 1.6% had a multiple pregnancy
- 58% of women having a singleton live birth delivered spontaneously, while deliveries using forceps accounted for 4% and vacuum extractions for 12%. 25% of women gave birth by Caesarean section, compared to 20% in 1999
- the average length of stay in hospital was 3.6 days, and was longer for women who had a Caesarean section (5.6 days)
- an estimated 1 in 5 births were to mothers from outside Ireland in 2007. This compares to 1 in 6 in 2004. In 2007, 7.2% of all births were to mothers from EU15 to EU27 Accession States, while mothers from Africa and Asia each accounted for 3.6% of all births.

## **SECTION ONE**

### **INTRODUCTION**



#### INTRODUCTION

Annual reports on perinatal statistics in Ireland are compiled from the statistical data collected in Part 3 of the standard Birth Notification Form (BNF01/2003) used to notify local registrars of all live births and stillbirths occurring in Ireland. This statistical data set, known as the National Perinatal Reporting System (NPRS), is a valuable resource and the only complete reporting system on births nationally. Since 1999 the Economic and Social Research Institute (ESRI) has been contracted by the Department of Health and Children, and more recently the Health Service Executive, to oversee the collection, processing, management, and reporting of data submitted to the NPRS. Previous reports in this series are available for download at http://www.esri.ie/health\_information/nprs/nprs\_reports/

#### **PURPOSE OF THIS REPORT**

The principal aim of this report is the presentation of national statistics on perinatal events for the year 2007. This data set includes all births from 22 weeks gestation to the end of the first week of life.

This report aims to

- describe the fundamental social and biological characteristics of mothers and their babies
- report on pregnancy outcomes with particular reference to perinatal mortality
- highlight important aspects of perinatal care
- stimulate a wider interest in research in the area of perinatal epidemiology
- promote a general recognition of the importance of having an accurate and complete perinatal reporting system at a national level.

#### STRUCTURE OF THIS REPORT

This chapter provides background information and details on data source, quality, and data elements.

**SECTION 2** contains summary data on selected variables since 1999 and trends in international comparisons.

**SECTION 3** contains tables relating to the following variables: maternal age, marital status, socio-economic group of mother and father (derived from collection of data on occupations), nationality of mother and father, maternal parity, distribution of previous stillbirths, interval since last birth, birthweight, gestational age, and month of birth.

SECTION 4 presents tables relating to variables which fall more naturally into the category of the use of health care facilities, and deals in particular with the following: type of antenatal care received, timing of first antenatal visit, length of mother's antenatal and postnatal stay, whether hospital admission was booked or not, mother's immunity to rubella, method of delivery, classification of outcome according to size of maternity unit, transfer of infants between hospitals, length of infant's hospital stay, and type of feeding.

SECTION 5 deals with perinatal outcomes in detail, examining perinatal deaths by birthweight and gestational age, with particular focus on cause of death. In this section tables are presented on the following variables: numbers of antepartum and intrapartum stillbirths and early neonatal deaths by birthweight groups and gestational age, numbers and rates for cause of death, age at death for early neonatal deaths, and numbers and percentages of stillbirths and early neonatal deaths undergoing post-mortem examinations.

APPENDICES include trends and tables on domiciliary births and source data tables for figures used throughout this report.

#### **CONVENTIONS USED IN THIS REPORT**

Singleton and multiple births are presented separately for each variable, since numbers and rates can vary considerably for the two groups. 1 Caution should be exercised when interpreting rates based on small numbers. Rates calculated from 20 observations or less are distinguished by italic type as a warning to the user that their reliability as a measure may be affected by the small number of observations.

Appendix I contains a guide cross referencing the tables in this report with those in *Perinatal* Statistics Report 2006.

Throughout this report, 'Not Stated' values are not included in the calculation of percentages used in discussion. 'Not Stated' values are shown in data tables. Where a particular value of a variable does not occur in a table, the row corresponding to that value is not included. This arises most frequently in relation to multiple births.

Unless otherwise stated, discussion on total births in this report includes live births and stillbirths for singleton and multiple births.

#### **NATIONAL MINIMUM DATA SET**

The NPRS data set provides for a national minimum data set, and is intended to serve as a basis for the development of a series of analyses overtime. Births are registered and notified on a standard four-part Birth Notification Form (BNF01/2003) used to notify local registrars of all live births and stillbirths occurring in Ireland. The third part of the form has all

<sup>&</sup>lt;sup>1</sup> Tables for singleton and multiple births are not presented in separate sections as in previous Perinatal Statistics Reports.

identifying information deleted and is sent to the HRID at the ESRI (see Appendix B). Figure 1.1 provides a diagrammatic representation of the information system around the registration of births. All the tables in this report are based on data contained in the third copy of the standard form.

The data collected by the NPRS can be grouped logically into demographic, clinical and administrative data as follows:

#### **DEMOGRAPHIC DATA**

- Date of infant's birth, sex of infant, multiple birth indicator
- Date of father's birth, father's county and country of residence, father's occupation, father's nationality (from 2004)
- Date of mother's birth, mother's county and country of residence, mother's occupation, mother's nationality (from 2004)
- Marital status of mother and date of present marriage

#### **CLINICAL DATA**

- Infant's birthweight and period of gestation, and mother's obstetric history (including parity)
- Autopsy indicator whether stillbirth occurred before or during labour
- Main cause of death and one secondary cause of death for infant
- Maternal immunity to rubella, method of delivery
- Infant's BCG vaccination
- Main maternal disease or condition and one secondary maternal disease or condition affecting the infant or foetus
- Main disease or congenital malformation and one secondary disease or congenital malformation affecting the infant or foetus

#### **ADMINISTRATIVE DATA**


- Hospital number, case number (unique to hospital of birth)
- Type of birth indicator, place of birth
- Date of last birth (live birth or stillbirth)
- Type of death indicator, place of death
- Type of antenatal care received, date of first visit to doctor and hospital during pregnancy
- Type of feeding, pre-booked admission indicator, dates of mother's admission and discharge, date of infant's discharge
- Infant transfer indicator, hospital number for hospital to which infant was transferred

The NPRS seeks to compile an accurate, complete and up-to-date data set on perinatal events over a specified time period. The national data sets on perinatal events are produced to the highest professional standards and comply with our commitments to confidentiality, access and timeliness.

The link between the birth notification form and the birth registration process for the General Register Office (GRO) ensures complete coverage of all births for the NPRS (see Figure 1.1). In addition, each birth is recorded using case reference numbers unique to each hospital/maternity unit. Missing records can be identified by using this unique case reference number system. Where a record is found to be missing, a request is made to each hospital/maternity unit to provide the birth notification form for any missing records. Similarly, independent midwives are requested to submit an annual summary of all births attended for a specified year, which is cross checked against the birth notification forms received.

The Information Technology and Support team at the HRID has developed a data entry and validation software system for NPRS data which is custom designed for our processing requirements. Data are received, either in electronic or paper format, from the hospitals/independent midwives and entered onto the NPRS data system. A comprehensive series of edits are built into this software system to check for completeness, validity, logical and coding errors at the point of data entry. The completeness of each record is dependent on those persons responsible for notification and registration of the births in each hospital/maternity unit throughout the country and each independent midwife working in the community.

FIGURE 1.1 NATIONAL PERINATAL REPORTING SYSTEM, DATA COLLECTION



Notes: CSO denotes Central Statistics Office. HSE denotes Health Service Executive.

## **SECTION TWO**

# TIME TRENDS AND INTERNATIONAL COMPARISONS



#### INTRODUCTION

This section presents data on selected statistics and three key indicators: perinatal mortality, birth rate and fertility. At a national level, data presented here show how these parameters have changed over time. International comparisons then allow the most recent statistics for Ireland to be compared with those for other European countries. The indicators for Ireland were estimated using data from the NPRS for the period 1999 to 2007.

Table 2.1 outlines trends in selected statistics for the period 1999 to 2007. Over the 1999 to 2007 period the average age of all mothers giving birth increased by just over one year from 30.06 years to 31.09 years. For single mothers, the average age increased from 24.74 years in 1999 to 27.06 years in 2007, while the percentage of total births to single mothers increased from 29.73 to 31.84 per cent. Average parity (number of previous live and still births) has declined from 1.10 to 1.00 over the period and the interval in years since delivery of the last birth has remained relatively stable over the period, ranging from 3.81 to 3.87 years. These data indicate that, on average, women giving birth in Ireland are doing so at an older age and having fewer births.

Trends in babies' characteristics have been largely consistent over the period 1999 to 2007. While the average birthweight for singleton births did not record any significant change from year to year, the average birthweight for multiple births varied somewhat. The average birthweight for multiple births ranged from a high of 2,509.85 grams in 2000 to a low of 2,421.36 grams in 2007. The average birthweight for all babies remained stable from one year to the next. The percentage of low birthweight babies showed little variation amongst singleton births but was more variable among multiple births. The percentage of total births recording low birthweights varied from 4.87 per cent in 2000 to 5.26 per cent in 2007.

Other statistics relating to the birth episode and post delivery are also outlined in Table 2.1. Booked admissions have consistently reported high percentages and have steadily increased to over 99 per cent for 2007. The percentages of women choosing combined antenatal care has also increased between 1999 and 2007 reaching a high of 79.01 per cent in 2006 and declining slightly in 2007 to 77.45 per cent. Method of delivery records declining trends for spontaneous and forceps deliveries for both singleton and multiple births, with total spontaneous deliveries falling from 64.46 per cent in 1999 to 57.26 per cent in 2007. The trend in delivery by caesarean section has increased from 20.43 per cent in 1999 to 26.24 per cent in 2007. (See also ESRI Working Paper 309, Recent Trends in the Caesarean Section Rate in Ireland, 1999-2006, Brick, A., and Layte, R., ESRI, August 2009.) Average lengths of stay for both mother and baby have declined between 1999 and 2007. Mother's average length of stay was recorded at 4.71 days in 1999 and has fallen steadily to 3.58 days in 2007. The decrease in average length of stay for babies has been less distinct, falling from 4.34 days in 1999 to 3.50 days in 2007. The percentage of women breastfeeding at discharge from hospital has risen from 36.16 per cent in 1999 to 45.22 per cent in 2007.

**TABLE 2.1**Trends in Selected Statistics, 1999-2007

Selected Statistics	1999	2000	2001	2002	2003	2004	2005	2006	2007
General Characteristics	1333	2000	2001	2002	2003	2007	2003	2000	2007
Average Age of Mother in years <sup>1</sup>	30.06	30.18	30.27	30.41	30.58	30.79	31.02	31.04	31.09
Average Maternal Parity <sup>1</sup>	1.10	1.09	1.07	1.06	1.05	1.05	1.05	1.02	1.00
Single Mothers (%) 1,2	29.73	30.09	30.14	30.01	30.20	30.63	30.71	31.53	31.84
Average Age of Single Mothers in years <sup>1,2</sup>	24.74	24.98	25.36	25.72	26.10	26.38	26.63	26.82	27.06
Average Number of Previous Stillbirths per 1,000 maternities <sup>1</sup>	14.03	13.06	13.57	12.43	12.02	12.60	12.79	10.57	10.76
Average Weight in grams <sup>3</sup>									
- Singleton Births	3,505.35	3,517.27	3,509.36	3,503.73	3,500.64	3,507.96	3,503.23	3,503.93	3,499.33
- Multiple Births	2,467.24	2,509.85	2,470.41	2,489.72	2,487.13	2,471.20	2,469.18	2,464.77	2,421.36
- Total Births	3,477.19	3,491.05	3,476.96	3,474.07	3,471.11	3,476.56	3,471.97	3,473.04	3,465.41
Low Birthweight <2500 grams (%) <sup>3</sup>									
- Singleton Births	3.80	3.84	3.84	3.76	3.87	3.71	3.76	3.74	3.79
- Multiple Births	47.71	43.35	46.46	44.18	45.07	45.56	45.62	46.38	50.44
- Total Births	4.99	4.87	5.17	4.94	5.07	4.98	5.02	5.01	5.26
Average Gestational Age at delivery in weeks <sup>1</sup>	39.48	39.46	39.47	39.45	39.44	39.38	39.42	39.40	39.38
Average Interval since last birth in years <sup>1</sup>	3.82	3.84	3.81	3.85	3.78	3.72	3.76	3.84	3.87
Perinatal Care									
Hospital & GP combined antenatal care (%) 1	72.53	72.43	73.29	74.07	75.86	77.29	77.47	79.01	77.45
Mother's Average Length of Stay in days <sup>4</sup>	4.71	4.54	4.38	4.09	3.95	3.85	3.84	3.72	3.58
Immunity to Rubella (%) <sup>1</sup>	89.63	87.76	86.84	86.49	87.30	90.49	90.26	92.21	92.36
Booked Hospital Admissions (%) 1	96.57	96.49	98.17	97.59	97.47	98.14	98.41	98.96	99.01
Mothers Breastfeeding (%) <sup>5</sup>	36.16	38.41	39.13	41.14	41.32	42.46	44.13	44.45	45.22
Post-mortems for Early Neonatal Deaths (%) <sup>6</sup>	44.37	39.41	33.55	42.07	41.72	38.16	24.43	26.85	30.77

**TABLE 2.1**Trends in Selected Statistics, 1999-2007 (contd.)

Selected Statistics	Selected Statistics		2000	2001	2002	2003	2004	2005	2006	2007
Perinatal Care										
Method of Delivery (%) <sup>7</sup>										
	Singleton Births									
	- Spontaneous	65.35	64.56	64.29	63.27	62.24	60.54	60.05	59.80	58.42
	- Caesarean	19.69	20.55	21.05	21.61	23.14	24.23	24.86	24.33	25.03
	- Forceps <sup>8</sup>	4.27	3.78	3.21	3.19	2.89	2.48	2.85	3.04	3.73
	Multiple Births									
	- Spontaneous	32.33	33.73	29.83	33.62	26.30	28.23	25.51	25.83	21.8
	- Caesarean	47.23	50.31	52.74	49.49	58.33	55.82	60.44	59.78	63.6
	- Forceps <sup>8</sup>	3.08	2.52	2.71	2.99	1.56	2.34	1.67	2.47	2.7
	Total Births									
	- Spontaneous	64.46	63.76	63.22	62.41	61.20	59.56	59.00	58.79	57.2
	- Caesarean	20.43	21.32	22.04	22.43	24.17	25.19	25.93	25.39	26.2
	- Forceps <sup>8</sup>	4.24	3.74	3.20	3.18	2.85	2.48	2.82	3.02	3.7
Infant's Average Length of Sta	0									
	- Singleton Births	4.17	4.08	3.89	3.68	3.62	3.55	3.52	3.47	3.2
	- Multiple Births	10.68	10.73	11.45	10.16	10.09	10.59	10.14	9.85	9.9
	- Total Births	4.34	4.25	4.12	3.87	3.80	3.76	3.72	3.66	3.50

Notes: 'Not Stated' or 'Not Known' values are not included in the calculation of percentages.

<sup>&</sup>lt;sup>2</sup> Data recording marital status as 'never married' are classified as 'single' marital status in this report.

<sup>&</sup>lt;sup>3</sup> Based on live births.

<sup>&</sup>lt;sup>4</sup> Based on maternities with live births.

<sup>&</sup>lt;sup>5</sup> Based on maternities with live births. Excludes early neonatal deaths.

<sup>&</sup>lt;sup>6</sup> Based on early neonatal deaths.

<sup>&</sup>lt;sup>7</sup> Based on live births. These figures were not based on total maternities but on total births in order to give a more accurate picture, as each infant in a multiple pregnancy can have a different method of delivery. See Tables 4.7a, 4.7b and 4.8 for data on all Methods of Delivery.

<sup>&</sup>lt;sup>8</sup> Excludes breech deliveries.

 $<sup>^{\</sup>rm 9}\,$  Based on live births. Excludes early neonatal deaths.

#### PERINATAL MORTALITY


In 2007 approximately 1 in every 180 births were stillborn and 1 in every 550 live births died in the first week of life. Despite improvements in reducing perinatal mortality in recent decades, the perinatal period continues to be a time of relatively high risk. Table 2.2 below shows the numbers and rates for total births and perinatal deaths for Ireland from 1999 to 2007. The perinatal mortality rate is the ratio of total stillbirths and early neonatal deaths to total live births and stillbirths, and is expressed as the rate per 1,000 live births and stillbirths. Stillbirths account for almost three quarters of perinatal deaths and are almost three times as common as early neonatal deaths. Set in this context, analysis of the foetal mortality rate is clearly a public health concern. Section 5 of this report focuses on perinatal outcomes in more detail, examining perinatal deaths by birthweight and gestational age, with particular focus on cause of death. In 1999 the perinatal mortality rate was 8.2 per 1,000 live births and stillbirths. This rate peaked at 8.9 per 1,000 in 2000 and declined to a low of 7.0 per 1,000 in 2006. For 2007 the perinatal mortality rate has increased to 7.3 per 1,000 live births and stillbirths (see Figure 2.1). For numbers and rates by singleton and multiple births see 'Singleton and Multiple Births' page 16.

**TABLE 2.2** Live Births, Stillbirths, Early Neonatal Deaths, and Perinatal Deaths, Ireland, 1999-2007

	1999	2000	2001	2002	2003	2004	2005	2006	2007			
Numbers												
Live Births	54,019	54,871	57,922	60,522	61,632	62,067	61,480	65,502	71,570			
Stillbirths	288	315	339	343	357	339	306	308	393			
Early Neonatal Deaths	157	176	160	170	177	168	139	152	132			
Perinatal Deaths	445	491	499	513	534	507	445	460	525			
Adjusted Perinatal Deaths	338	355	356	396	384	382	330	345	383			
					Rates							
Stillbirths	5.3	5.7	5.8	5.6	5.8	5.4	5.0	4.7	5.5			
Early Neonatal Deaths	2.9	3.2	2.8	2.8	2.9	2.7	2.3	2.3	1.8			
Perinatal Deaths	8.2	8.9	8.6	8.4	8.6	8.1	7.2	7.0	7.3			
Adjusted Perinatal Deaths	6.2	6.4	6.1	6.5	6.2	6.1	5.4	5.3	5.3			

Note: See Appendix A for further details on calculation of rates.


FIGURE 2.1 Stillbirth, Early Neonatal and Perinatal Mortality Rates, Ireland, 1999-2007



Note: See Table 2.2 for data.

Figure 2.2 presents the 2007 perinatal mortality rate for 20 European countries. In 2007, Ireland had a comparatively high perinatal mortality rate, 7.3 per 1,000 live births and stillbirths, ranking 13th out of 20 countries included in the Eurostat system. Of these countries, Bulgaria recorded the highest perinatal mortality rate in 2007, at 11.0 per 1,000 live births and stillbirths. The country with the lowest perinatal mortality rate was the Czech Republic at 4.0 per 1,000 live births and stillbirths. When countries are compared it should be remembered that rates are affected by many factors such as national policies on abortion, screening for congenital anomalies, availability of services related to fertility treatment, differences in the comprehensiveness and accuracy of registration and notification systems, and variations in definitions employed (in particular, for stillbirth).

FIGURE 2.2 Perinatal Mortality Rates for Selected EU Countries, 2007



'Population and Social Conditions', Eurostat, Data Navigation Tree, date consulted August 2009. http://epp.eurostat.ec.europa.eu Source:

- \* National Figure compiled by NPRS. See Table 2.2 for data.
- \*\* Health Statistics Quarterly 42, 66. Available on the Office for National Statistics website at: http://www.statistics.gov.uk/downloads/theme\_health/HSQ42.pdf

#### **BIRTH RATES<sup>2</sup>**


The birth rate is based on the total number of live births per 1,000 population.<sup>3</sup> Figure 2.3 shows trends in birth rates for selected EU countries since 1999. For the countries reviewed, Ireland has consistently had the highest birth rate over the period. While Ireland's birth rate was closer to that of other European countries in 1999 and 2000, since then the gap has grown between Ireland and France, the European country closest to Ireland in terms of birth rates. Ireland's birth rate peaked at 15.5 per 1,000 population in 2002 and 2003, declining slightly in 2004 and 2005 and then returning to 15.4 per 1,000 in 2006. In 2007, the birth rate increased substantially to 16.5 per 1,000 population which is the highest recorded rate for the 27 EU countries presented in Figure 2.4.4 After Ireland, the country with the next highest birth rate is France (12.9) followed by the United Kingdom (12.7). In 2007, Germany (8.3) had the lowest birth rate with Austria (9.2) next in line.

<sup>&</sup>lt;sup>2</sup> It should be noted that the crude birth rate could rise or fall depending on the numbers of women in the age cohorts from 15 to 49 years old. Changes in the crude birth rate often mask underlying trends in fertility. The next section is, therefore, concerned with changes in fertility over time.

The population estimates used in the denominator of the birth rate were calculated by the Department of Health and Children and obtained from Ireland and Northern Ireland's Population Health Observatory (INIsPHO) PHIS (Online) Indicators System, http://www.inispho.org/phis/indicators, date consulted August 2009. For the Census year 2002, population data from PHIS concur with the Census. Population estimates for 2006 were obtained from Census 2006. Population estimates for 2007 were obtained from the Economic and Social Research Institute.

Provisional data obtained from the Vital Statistics section, CSO estimates the birth rate to be 16.5 for 2007 (personal communication, October 2009).

FIGURE 2.3 Trends in Birth Rates for Selected EU Countries, 1999-2007




Source:

'Population and Social Conditions', Eurostat, Data Navigation Tree, date consulted August 2009. http://epp.eurostat.ec.europa.eu

\* Data for Ireland are based on NPRS data on live births from 1999 onwards. Population estimates were obtained from Ireland and Northern Ireland's Population Health Observatory (INIsPHO) PHIS (Online) Indicators System, http://www.inispho.org/phis/indicators, date consulted August 2009. For the Census year 2002, population data from PHIS concur with the Census. Population estimates for 2006 were obtained from Census 2006. Population estimates for 2007 were obtained from the Economic and Social Research Institute.

FIGURE 2.4 Birth Rates per 1,000 Population for EU Countries, 2007



Source:


'Population and Social Conditions', Eurostat, Data Navigation Tree, date consulted August 2009. http://epp.eurostat.ec.europa.eu

\* Data for Ireland are based on NPRS data on live births. Population estimates for 2007 were obtained from the Economic and Social Research Institute.

#### **FERTILITY**

Figure 2.5 shows the trend in the total period fertility rate (TPFR) in Ireland from 1999 to 2007. The TPFR for a given year indicates the number of children a woman could expect to have if the age-specific fertility rates (ASFR) for that year applied throughout her fertile years. The Irish TPFR has increased overall from an average of 1.91 in 1999 to 2.06 in 2007, the highest since the early ninties. 6 The ASFRs used in the calculation of the TPFR, and also shown in Figure 2.5, indicate differences in fertility patterns between younger and older age groups. Compared to 1999, only the age group categories 15 to 19 years and 25 to 29 years recorded lower ASFRs in 2007; all other age group categories recorded higher ASFRs in 2007, some more than others. Women in the older age group categories (35 to 49 years) recorded increased rates each year over the period 1999 to 2007, whilst women in the younger age group categories (15 to 29 years) recorded decreasing rates each year up to 2005, with the 20 to 29 year age groups beginning to increase in 2006. The most significant change noted is for the increase in the ASFR for the 35 to 39 year age group, which in 2007 remains above the 25 to 29 year age group for a third year. Since 2005 the number of live births per 1,000 women has increased for all age group categories, with marked increases for 2007 in particular. In all years the 30 to 34 year age group recorded the highest ASFR. Notably, the ASFR for 40 to 44 year age group remained greater than that reported for the 15 to 19 year age group in 2007, a trend also observed in 2005 and 2006.

FIGURE 2.5 Age-Specific Fertility Rates (ASFRs) and Total Period Fertility Rates (TPFRs), Ireland, 1999-2007




Total period fertility rates (TPFRs) and age-specific fertility rates (ASFRs) are calculated using NPRS data from 1999 Source: onwards. Female population estimates for the years 1999 to 2005 were obtained from Ireland and Northern Ireland's Population Health Observatory (INISPHO) PHIS (Online) Indicators System, http://www.inispho.org/phis/indicators, date consulted August 2009. For the Census year 2002, population data from PHIS concur with the Census. Population estimates for 2006 were obtained from Census 2006. Population estimates for 2007 were obtained from the Economic and Social Research Institute.

<sup>&</sup>lt;sup>5</sup> ASFR is the number of live births to women in a particular age group divided by the female population in that age group, multiplied by 1,000. The TPFR is compiled by summing the ASFRs for the relevant five-year age groups, dividing by 1,000 and

<sup>&</sup>lt;sup>6</sup> Report on Vital Statistics 2006, Central Statistics Office, Ireland

At 2.06, for the first time since 1999 the TPFR for Ireland has almost reached the level of 2.1 required for the long-term replacement of the population in the absence of any net inward migration. Using data from Eurostat and the NPRS, Figure 2.6 shows that, for 2007, Ireland had the highest TPFR in the EU at 2.06, with France just behind with a TPFR of 1.98. For 2007, Slovakia had the lowest TPFR (1.25) of the EU countries for which data were available.

FIGURE 2.6 Total Period Fertility Rates for Selected EU Countries, 2007



Source: 'Population and Social Conditions', Eurostat, Data Navigation Tree, date consulted August 2009, http://epp.eurostat.ec.europa.eu

Population estimates for 2007 were obtained from the Economic and Social Research Institute.

#### SINGLETON AND MULTIPLE BIRTHS

Table 2.3 below shows the numbers of singleton and multiple births each year for the period 1999 to 2007. In 2007 there were 69,730 singleton births, 2,192 twin births, 87 triplets, and 4 quadruplets reported to NPRS. Of these, 50 births weighed less than 500 grams: 38 singleton births and 12 multiple births. In accordance with the World Health Organization (WHO) guidelines the National Perinatal data set includes only births weighing 500 grams or more. For the purposes of this report, therefore, there were 69,692 singleton births, 2,185 twin births, 83 triplet births and 3 quadruplet births in 2007. The twinning rate was 15.4 per 1,000 maternities in 2007. Due to the small number of multiple births the twinning rate can vary from year to year. During 1999 to 2007 the average twinning rate was 14.7 per 1,000 maternities.

<sup>\*</sup> Data for Ireland are based on NPRS data on live births.

**TABLE 2.3** Singleton and Multiple Births, 1999-2007

	1999	2000	2001	2002	2003	2004	2005	2006	2007
Singleton births	52,830	53,737	56,422	59,071	60,164	60,499	59,902	63,839	69,692
Twin births	1,392	1,397	1,756	1,728	1,747	1,849	1,833	1,916	2,185
Triplet births	85	48	71	54	78	54	47	51	83
Quadruplet births	0	4	7	12	0	4	4	4	3
Quintuplet births	0	0	5	0	0	0	0	0	0
Twinning rate (per 1,000 maternities)	14.8	12.8	15.3	14.4	14.3	15.0	15.1	14.8	15.4

Note: See Appendix A for details on calculation of twinning rate.

The number of stillbirths and early neonatal deaths for singleton and multiple births between 1999 and 2007 is presented in Table 2.4. In all years higher mortality rates are estimated for multiple births when compared to singleton births, though the number for multiple births was small. In 2007, the stillbirth mortality rate for singleton births was 5.4 per 1,000 live births and stillbirths. The early neonatal mortality rate for singleton births was 1.7 per 1,000 live births.

**TABLE 2.4** Live Births, Stillbirths and Early Neonatal Deaths, Numbers and Rates, for Singleton and Multiple Births, 1999-2007

	1999	2000	2001	2002	2003	2004	2005	2006	2007
					Numbers				
Total Births	54,307	55,186	58,261	60,865	61,989	62,406	61,786	65,810	71,963
Singleton	52,830	53,737	56,422	59,071	60,164	60,499	59,902	63,839	69,692
Multiple	1,477	1,449	1,839	1,794	1,825	1,907	1,884	1,971	2,271
Live Births	54,019	54,871	57,922	60,522	61,632	62,067	61,480	65,502	71,570
Singleton	52,556	53,442	56,115	58,752	59,837	60,186	59,621	63,555	69,318
Multiple	1,463	1,429	1,807	1,770	1,795	1,881	1,859	1,947	2,252
Stillbirths	288	315	339	343	357	339	306	308	393
Singleton	274	295	307	319	327	313	281	284	374
Multiple	14	20	32	24	30	26	25	24	19
<b>Early Neonatal</b>	157	176	160	170	177	168	139	152	132
Deaths	_		_						
Singleton	130	153	136	149	156	140	109	129	118
Multiple	27	23	24	21	21	28	30	23	14
	I				Rates				
Stillbirths	5.3	5.7	5.8	5.6	5.8	5.4	5.0	4.7	5.5
Singleton	5.2	5.5	5.4	5.4	5.4	5.2	4.7	4.4	5.4
Multiple	9.5	13.8	17.4	13.4	16.4	13.6	13.3	12.2	8.4
Early Neonatal	2.9	3.2	2.8	2.8	2.9	2.7	2.3	2.3	1.8
Deaths	2.5	2.0	2.4	2 -	2.0	2.2	4.0	2.0	4 -
Singleton	2.5	2.9	2.4	2.5	2.6	2.3	1.8	2.0	1.7
Multiple	18.5	16.1	13.3	11.9	11.7	14.9	16.1	11.8	6.2

Note: See Appendix A for details on calculation of rates.

## **SECTION THREE**

### **GENERAL CHARACTERISTICS**



#### **INTRODUCTION**

This section presents more detailed analyses of data reported to the NPRS for 2007. Singleton and multiple births are reported in separate tables. For selected variables, the following tables outline data for live births, stillbirths and early neonatal deaths, together with their corresponding rates. The perinatal mortality rate and adjusted perinatal mortality rate are also outlined in each table. Tables and figures are presented in some detail for a number of variables relating to the mother (age, marital status, occupation, nationality, parity, number of previous stillbirths, and interval since last birth), father (occupation and nationality), and infant (birthweight, gestational age, and month of birth).

#### **MATERNAL AGE**

The average maternal age for all births rose from 30.6 to 31.1 years over the five years 2003 to 2007. The number of total births to mothers aged 19 years and under decreased from 2,880 in 2003 to 2,521 in 2007 (see Tables 3.1a and 3.1b). The percentage of births to mothers aged 19 years and under fell from 4.6 to 3.5 per cent over the five-year period. The number of births to mothers 35 years or over increased from 14,540 in 2003 to 19,128 in 2007 (see Tables 3.1a and 3.1b), representing an increase from 23.5 to 26.6 per cent of births.

**TABLE 3.1a: SINGLETON BIRTHS** 

Age of Mother

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Age of Mother	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Under 15 Years	10	0.0	0	0.0	0	0.0	0.0	0.0
15-19 Years	2,428	3.5	14	5.7	7	2.9	8.6	5.7
20-24 Years	8,814	12.7	54	6.1	14	1.6	7.7	6.2
25-29 Years	16,258	23.5	82	5.0	22	1.4	6.4	4.9
30-34 Years	23,652	34.1	100	4.2	42	1.8	6.0	4.1
35-39 Years	15,179	21.9	97	6.3	30	2.0	8.3	5.6
40-44 Years	2,858	4.1	20	6.9	3	1.0	8.0	5.9
45 Years and Over	103	0.1	1	9.6	0	0.0	9.6	0.0
Not Stated	16	0.0	6	272.7	0	0.0	272.7	272.7
Total	69,318	100	374	5.4	118	1.7	7.1	5.1

**TABLE 3.1b: MULTIPLE BIRTHS** 

Age of Mother

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007


Age of Mother	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
15-19 Years	69	3.1	0	0.0	1	14.5	14.5	14.5
20-24 Years	125	5.6	1	7.9	4	32.0	39.7	39.7
25-29 Years	359	15.9	1	2.8	0	0.0	2.8	2.8
30-34 Years	841	37.3	5	5.9	6	7.1	13.0	9.5
35-39 Years	707	31.4	12	16.7	3	4.2	20.9	19.5
40-44 Years	140	6.2	0	0.0	0	0.0	0.0	0.0
45 Years and Over	11	0.5	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

<sup>&</sup>lt;sup>7</sup> For data on 2003, see *Report on Perinatal Statistics for 2003*, The Economic and Social Research Institute, December 2006.

<sup>&</sup>lt;sup>8</sup> Throughout this section, 'Not Stated' values are not included in the calculation of percentages for discussion.

When we examine perinatal mortality rates by mother's age, the importance of focusing on both early and late childbearing becomes apparent. As highlighted in Figure 3.1, perinatal mortality rates are highest for mothers aged 19 years and under and 35 years and over, with a rate of 8.7 per 1,000 live births and stillbirths for both categories. The lowest stillbirth and total perinatal mortality rates were recorded in the 30-34 year age group (4.3 and 6.2 per 1,000 live births and stillbirths respectively). Rates for early neonatal mortality were fairly consistent for mothers aged over 20 years, with slightly higher rates recorded for mothers aged 19 years and under.

FIGURE 3.1 Age-Specific Stillbirth, Early Neonatal and Perinatal Mortality Rates, Total Births, 2007



Note: See Appendix H, Table H1 for data.

#### **MOTHER'S MARITAL STATUS**

The distribution of total births by mother's marital status has remained fairly consistent in recent years with over two-thirds to married women, and just under one third to single women (see Figure 3.2). In the case of multiple births, these proportions differ; just over 74 per cent of multiple births were to married women (see Table 3.2b). Although a very small proportion of total births, the percentage of births to mothers whose marital status is 'divorced' is increasing, and has doubled from 0.3 per cent of total births in 2003 to 0.6 per cent of total births in 2007 (see Figure 3.2). The distribution of single mothers by mother's area of residence is shown in Figure 3.3. Dublin City recorded the highest percentage of single mothers at just over 40 per cent, and was followed by counties Louth, Wexford and Tipperary South recording between 36 and 40 per cent. Galway recorded the lowest number of single mothers at 24.2 per cent and was closely followed by Roscommon at 24.3 per cent.10

<sup>&</sup>lt;sup>9</sup> Data recording marital status as 'never married' are classified as 'single' marital status in discussion in this report.

<sup>&</sup>lt;sup>10</sup> Percentages of single mothers by county will be affected by the female age populations for each county.

**TABLE 3.2a: SINGLETON BIRTHS** 

Mother's Marital Status

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Mother's Marital Status	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Married	45,897	66.2	220	4.8	77	1.7	6.4	4.3
Never Married	22,142	31.9	143	6.4	40	1.8	8.2	6.7
Widowed	87	0.1	0	0.0	0	0.0	0.0	0.0
Separated	789	1.1	9	11.3	1	1.3	12.5	8.8
Divorced	399	0.6	1	2.5	0	0.0	2.5	2.5
Not Stated	4	0.0	1	200.0	0	0.0	200.0	200.0
Total	69,318	100	374	5.4	118	1.7	7.1	5.1

#### **TABLE 3.2b: MULTIPLE BIRTHS**


Mother's Marital Status

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Mother's Marital Status	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Married	1,668	74.1	15	8.9	9	5.4	14.3	11.9
Never Married	531	23.6	3	5.6	5	9.4	15.0	15.0
Widowed	2	0.1	0	0.0	0	0.0	0.0	0.0
Separated	28	1.2	1	34.5	0	0.0	34.5	34.5
Divorced	22	1.0	0	0.0	0	0.0	0.0	0.0
Not Stated	1	0.0	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8


Notes:

FIGURE 3.2 Percentage of Births by Mother's Marital Status, Total Births, 2003-2007



See Appendix H, Table H2 for data. 'Not Stated' values are not included in the calculation of percentages. Note:

FIGURE 3.3 Percentage of Births to Single Mothers by Mother's County of Residence, Total Births, 2007



See Appendix H, Table H3 for data. 'Not Stated' values are not included in the calculation of percentages. Example: Of all births to mothers resident in Carlow, 35.7% were to single mothers.

# **OCCUPATION**

### **MOTHER'S OCCUPATION**

Almost half of total births were to mothers whose occupation was recorded as either 'home duties' (23.6 per cent) or 'intermediate non-manual workers' (22.9 per cent). The occupations with the next highest percentage of births were 'lower professional', reported as 13.9 per cent of total births and 'other non-manual workers', reported as 12.5 per cent of total births (see Tables 3.3a and 3.3b). Where rates are calculated on more than 20 observations, the occupations recording the highest perinatal mortality rates were 'unemployed' and 'home duties' (see Figure 3.4).

**TABLE 3.3a: SINGLETON BIRTHS** 

Mother's Occupation

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Mother's Occupation	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Farmers & Farm Managers	65	0.1	2	29.9	0	0.0	29.9	29.9
Other Agricultural Occupations & Fishermen	89	0.1	0	0.0	0	0.0	0.0	0.0
Higher Professional	4,168	6.0	21	5.0	7	1.7	6.7	4.1
Lower Professional	9,540	13.8	40	4.2	17	1.8	5.9	4.1
Employers & Managers	4,954	7.1	21	4.2	9	1.8	6.0	3.6
Salaried Employees	1,040	1.5	7	6.7	3	2.9	9.6	5.8
Intermediate Non-Manual Workers	15,806	22.8	59	3.7	21	1.3	5.0	3.7
Other Non- Manual Workers	8,667	12.5	41	4.7	9	1.0	5.7	4.4
Skilled Manual Workers	1,238	1.8	5	4.0	2	1.6	5.6	4.8
Semi-Skilled Manual Workers	2,385	3.4	15	6.3	4	1.7	7.9	5.8
Unskilled Manual Workers	7	0.0	0	0.0	0	0.0	0.0	0.0
Unemployed	2,306	3.3	27	11.6	4	1.7	13.3	10.7
Not Classifiable	2,595	3.7	20	7.6	5	1.9	9.6	7.7
Home Duties	16,315	23.5	111	6.8	36	2.2	8.9	6.5
Not Stated	143	0.2	5	33.8	1	7.0	40.5	34.0
Total	69,318	100	374	5.4	118	1.7	7.1	5.1

*Note*: See Appendix C for a description of the classification system for occupation.

**TABLE 3.3b: MULTIPLE BIRTHS** 


Mother's Occupation

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Mother's Occupation	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Other Agricultural Occupations & Fishermen	4	0.2	0	0.0	0	0.0	0.0	0.0
Higher Professional	155	6.9	2	12.7	0	0.0	12.7	6.4
Lower Professional	365	16.2	3	8.2	0	0.0	8.2	8.2
Employers & Managers	167	7.4	1	6.0	4	24.0	29.8	24.0
Salaried Employees	45	2.0	0	0.0	0	0.0	0.0	0.0
Intermediate Non-Manual Workers	565	25.1	3	5.3	2	3.5	8.8	7.1
Other Non- Manual Workers	237	10.5	1	4.2	0	0.0	4.2	4.2
Skilled Manual Workers	45	2.0	0	0.0	1	22.2	22.2	22.2
Semi-Skilled Manual Workers	63	2.8	1	15.6	0	0.0	15.6	15.6
Unemployed	48	2.1	0	0.0	0	0.0	0.0	0.0
Not Classifiable	71	3.2	3	40.5	3	42.3	81.1	81.1
Home Duties	480	21.3	5	10.3	4	8.3	18.6	16.5
Not Stated	7	0.3	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

Note: See note to Table 3.3a.

FIGURE 3.4 Perinatal Mortality Rates by Mother's Occupation, Total Births, 2007



Note: See Appendix H, Table H4 for data. Rates based on 20 or less observations are not included in this Figure.

### FATHER'S OCCUPATION<sup>11</sup>

For over one-quarter of total births, the father's occupation was recorded as 'skilled manual workers'. Apart from 'not applicable', the next highest occupational grouping, 'other nonmanual workers', was recorded as 9.6 per cent of total births (see Tables 3.4a and 3.4b). Where rates are calculated on more than 20 observations, the occupations recording the highest perinatal mortality rates were amongst the lower socio-economic groups, where father's occupations were recorded as either 'unemployed' or 'semi-skilled manual workers' (see Figure 3.5).

The similar pattern observed among perinatal mortality rates for both mother and father's occupation indicate that socio economic grouping may have a bearing on perinatal outcomes. For further discussion and analysis see Working Paper 312, Layte, R., and Clyne, B., Did the Celtic Tiger Decrease Socio-Economic Differentials in Perinatal Mortality in Ireland? ESRI, September 2009.

<sup>&</sup>lt;sup>11</sup> The occupation category 'Not applicable' applies to father's occupation only. Where mother's marital status is recorded as 'never married', 'widowed', 'separated' or 'divorced', the GRO does not require father's details to register the birth. This information, therefore, is generally not provided in these circumstances. Where father's details are provided, they are included in the statistics.

**TABLE 3.4a: SINGLETON BIRTHS** 

Father's Occupation

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Father's Occupation	Live Births	Per Cent	Stillbirths	Stillbirth Rate	Early Neonatal	Early Neonatal	Perinatal Mortality	Adjusted PMR
		(%) Live Births			Deaths	Mortality Rate	Rate	
Farmers & Farm Managers	2,101	3.0	9	4.3	5	2.4	6.6	4.3
Other Agricultural Occupations & Fishermen	712	1.0	4	5.6	0	0.0	5.6	5.6
Higher Professional	5,314	7.7	22	4.1	6	1.1	5.2	3.2
Lower Professional	3,280	4.7	17	5.2	5	1.5	6.7	4.9
Employers & Managers	5,710	8.2	21	3.7	7	1.2	4.9	3.5
Salaried Employees	1,506	2.2	5	3.3	1	0.7	4.0	2.7
Intermediate Non-Manual Workers	5,471	7.9	27	4.9	9	1.6	6.5	4.9
Other Non- Manual Workers	6,590	9.5	32	4.8	9	1.4	6.2	4.4
Skilled Manual Workers	17,416	25.1	59	3.4	27	1.6	4.9	3.4
Semi-Skilled Manual Workers	3,106	4.5	14	4.5	8	2.6	7.1	4.2
Unskilled Manual Workers	822	1.2	4	4.8	2	2.4	7.3	2.4
Unemployed	2,107	3.0	17	8.0	5	2.4	10.4	6.6
Not Classifiable	1,544	2.2	8	5.2	2	1.3	6.4	5.8
Not Applicable	13,182	19.0	120	9.0	31	2.4	11.4	9.2
Home Duties	71	0.1	1	13.9	0	0.0	13.9	0.0
Not Stated Total	386	0.6 <b>100</b>	14 <b>374</b>	35.0 <b>5.4</b>	1 <b>118</b>	2.6 <b>1.7</b>	37.5 <b>7.1</b>	22.8 <b>5.1</b>
iotai	69,318	100	3/4	5.4	118	1./	7.1	5.1

Note: See Appendix C for a description of the classification system for occupation.

**TABLE 3.4b: MULTIPLE BIRTHS** 


Father's Occupation

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Father's Occupation	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Farmers & Farm Managers	62	2.8	0	0.0	0	0.0	0.0	0.0
Other Agricultural Occupations & Fishermen	16	0.7	0	0.0	0	0.0	0.0	0.0
Higher Professional	193	8.6	3	15.3	0	0.0	15.3	10.3
Lower Professional	107	4.8	0	0.0	0	0.0	0.0	0.0
Employers & Managers	222	9.9	2	8.9	3	13.5	22.3	9.0
Salaried Employees	72	3.2	0	0.0	0	0.0	0.0	0.0
Intermediate Non-Manual Workers	222	9.9	2	8.9	0	0.0	8.9	8.9
Other Non- Manual Workers	219	9.7	0	0.0	0	0.0	0.0	0.0
Skilled Manual Workers	517	23.0	5	9.6	4	7.7	17.2	17.2
Semi-Skilled Manual Workers	105	4.7	1	9.4	0	0.0	9.4	9.4
Unskilled Manual Workers	21	0.9	0	0.0	0	0.0	0.0	0.0
Unemployed	69	3.1	0	0.0	0	0.0	0.0	0.0
Not Classifiable	38	1.7	0	0.0	0	0.0	0.0	0.0
Not Applicable	360	16.0	3	8.3	5	13.9	22.0	22.0
Home Duties	5	0.2	0	0.0	0	0.0	0.0	0.0
Not Stated	24	1.1	3	111.1	2	83.3	185.2	185.2
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

Note: See note to Table 3.4a.

FIGURE 3.5 Perinatal Mortality Rates by Father's Occupation, Total Births, 2007



Note: See Appendix H, Table H5 for data. Rates based on 20 or less observations are not included in this Figure.

### **NATIONALITY**

### **MOTHER'S NATIONALITY**

Information on mother's nationality was reported for the first time in 2004. For the purposes of this report mother's nationality is recorded as 'place of birth'. See Appendix F for a detailed description on classification of nationality. In 2004, 84.4 per cent of total births were to women born in Ireland; 12 this figure fell to 79.3 per cent in 2007 (see Tables 3.5a and 3.5b). Babies born to mothers from the EU15 to EU27 accession states accounted for 7.2 per cent of births in 2007. 13 Figure 3.6 shows the percentage of total births by nationality of mothers accounting for the majority of total births to non-Irish mothers in 2007. Between 2004 and 2007 the percentage of total births to mothers within the EU15 to EU27 accession states category increased from less than 1 per cent to over 7 per cent whilst births to mothers from Africa decreased from 5.6 per cent to 3.6 per cent. Births to mothers from the UK remained relatively stable over this time period and births to mothers from Asia increased from 2.5 per cent to 3.6 per cent of total births (see Figure 3.6).

In 2007, higher mortality rates were recorded for babies born to mothers from Africa (10.5 per 1,000 live births and stillbirths) when compared to births to Irish mothers at 7.1 per 1,000 live births and stillbirths. Lower mortality rates were reported, however, for babies born to mothers from the EU15 to EU27 accession states, 5.7 per 1,000 live births and stillbirths (see Tables 3.5a and 3.5b).

The average maternal age for total births in 2007 is 31.1 years and, as demonstrated in Figure 3.7, there is some variation when mother's age is examined across nationality groups. Mothers from the EU15 to EU27 accession group are on average younger at 27.0 years, as were mothers from the rest of Europe (29.3 years). Mother's from the EU15 (excluding Ireland and UK) and Australia recorded the highest average maternal age at 32.5 years and 32.4 years respectively. For most other nationalities, the average age was closer to the average age for total births.

<sup>&</sup>lt;sup>12</sup> For data on 2004, see *Report on Perinatal Statistics for 2004*, The Economic and Social Research Institute, December 2007.

<sup>&</sup>lt;sup>13</sup> In previous reports, prior to the accession of Bulgaria and Romania to the EU, this category was titled EU15 to EU25 accession states. On 1 January 2007, Bulgaria and Romania joined the European Union, giving 27 Member States. In 2007, less than 1 per cent of total births were to mothers whose nationality was recorded as either Bulgarian or Romanian.

**TABLE 3.5a: SINGLETON BIRTHS** 

Mother's Nationality

Notes:

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Mother's Nationality	Live Births	Per Cent		t (%) Live B r's Marital	•	Stillbirths	Per Cent (%)		t (%) Stillbi r's Marital		Stillbirth Rate	Early Neonatal	Early Neonatal	Perinatal Mortality	Adjusted PMR
		(%) Live Births	Married	Never Married	Other*		Stillbirths	Married	Never Married	Other*		Deaths	Mortality Rate	Rate	
Ireland	54,714	78.9	52.4	25.4	1.2	297	79.4	44.9	31.8	2.7	5.4	91	1.7	7.1	5.0
UK	2,053	3.0	1.8	1.1	0.1	7	1.9	1.6	0.3	0.0	3.4	4	1.9	5.3	3.9
EU15 (excluding Ireland & UK)	931	1.3	0.8	0.5	0.0	1	0.3	0.0	0.3	0.0	1.1	1	1.1	2.1	2.1
EU15 to EU27 Accession States	5,022	7.2	4.0	3.0	0.3	25	6.7	3.7	2.7	0.3	5.0	4	0.8	5.7	4.8
Rest of Europe	642	0.9	0.7	0.2	0.0	2	0.5	0.5	0.0	0.0	3.1	2	3.1	6.2	4.7
Africa	2,470	3.6	2.5	0.9	0.1	18	4.8	3.2	1.6	0.0	7.2	8	3.2	10.5	8.1
Asia	2,546	3.7	3.0	0.6	0.1	13	3.5	2.9	0.5	0.0	5.1	5	2.0	7.0	5.5
America	639	0.9	0.7	0.2	0.0	4	1.1	0.8	0.3	0.0	6.2	1	1.6	7.8	7.8
Australia	100	0.1	0.1	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
New Zealand and Other Oceania	40	0.1	0.0	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Other Nationality	12	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Not Stated	149	0.2	0.1	0.1	0.0	7	1.9	1.1	0.8	0.0	44.9	2	13.4	57.7	39.2
Total	69,318	100	66.2	31.9	1.8	374	100	58.8	38.2	2.9	5.4	118	1.7	7.1	5.1


\* Includes; divorced, married but separated, widowed, and not stated. See Appendix F for a description of the classification system used for nationality. Mother's Nationality

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Mother's Nationality	Live Births	Per Cent		t (%) Live B r's Marital	•	Stillbirths	Per Cent (%)		t (%) Stillbi r's Marital		Stillbirth Rate	Early Neonatal	Early Neonatal	Perinatal Mortality	Adjusted PMR
		(%) Live Births	Married	Never Married	Other*		Stillbirths	Married	Never Married	Other*		Deaths	Mortality Rate	Rate	
Ireland	1,877	83.3	63.2	18.6	1.6	16	84.2	68.4	10.5	5.3	8.5	13	6.9	15.3	13.2
UK	58	2.6	1.6	1.0	0.0	1	5.3	5.3	0.0	0.0	16.9	0	0.0	16.9	16.9
EU15 (excluding Ireland & UK)	34	1.5	0.9	0.6	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
EU15 to EU27 Accession States	107	4.8	2.3	1.9	0.5	1	5.3	0.0	5.3	0.0	9.3	1	9.3	18.5	18.5
Rest of Europe	16	0.7	0.7	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Africa	95	4.2	3.0	1.1	0.1	1	5.3	5.3	0.0	0.0	10.4	0	0.0	10.4	10.4
Asia	41	1.8	1.6	0.3	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
America	14	0.6	0.5	0.1	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Australia	6	0.3	0.3	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Not Stated	4	0.2	0.0	0.1	0.1	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Total	2,252	100	74.1	23.6	2.4	19	100	78.9	15.8	5.3	8.4	14	6.2	14.5	12.8

Notes: \* Includes; divorced, married but separated, widowed, and not stated. See note to Table 3.5a.


FIGURE 3.6 Percentage of Births by Nationality for Non-Irish Mothers, Total Births, 2004-2007\*



See Appendix H, Table H6 for data. 'Not Stated' values are not included in the calculation of percentages. Notes: On 1 January 2007, Bulgaria and Romania joined the European Union, giving 27 Member States. Prior to 2007 there were 25 Member States in the European Union. In 2007, less than 1 per cent of total births were to mothers whose nationality was recorded as either Bulgarian or Romanian.

\*Mothers from the UK, EU15 to EU27 Accession States, Africa, and Asia accounted for 17.4 per cent of total births in 2007.

FIGURE 3.7 Average Age of Mother by Mother's Nationality, Total Births, 2007



Note: See Appendix H, Table H7 for data. Information on father's nationality (place of birth) was also reported for the first time in 2004 and 2007 data is shown in Tables 3.6a and 3.6b. See Appendix F for a detailed description on classification of nationality. Father's nationality was recorded as 'not applicable' for over 23 per cent of all births, decreasing from 28.4 per cent in 2004. Excluding these births, the largest proportion, 79.8 per cent of total births, occurred to Irish fathers, compared to 84.4 per cent in 2004. Father's nationality shows a similar pattern to mother's nationality, with the next highest proportions of births occurring to fathers from EU15 to EU27 accession states (6 per cent), followed by fathers from Africa (3.9 per cent).

<sup>&</sup>lt;sup>14</sup> The nationality category 'Not applicable' applies under father's nationality only. Where mother's marital status is recorded as 'never married', 'widowed', 'separated' or 'divorced', the GRO does not require father's details to register the birth. This information, therefore, is generally not provided in these circumstances. Where father's details are provided, they are included in the statistics.

For data on 2004, see Report on Perinatal Statistics for 2004, The Economic and Social Research Institute, December 2007.

**TABLE 3.6a: SINGLETON BIRTHS** 

Father's Nationality

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Father's Nationality	Live Births	Per Cent		t (%) Live B r's Marital :		Stillbirths	Per Cent (%)		t (%) Stillbi r's Marital		Stillbirth Rate	Early Neonatal	Early Neonatal	Perinatal Mortality	Adjusted PMR
		(%) Live Births	Married	Never Married	Other*		Stillbirths	Married	Never Married	Other*		Deaths	Mortality Rate	Rate	
Ireland	41,853	60.4	52.1	7.8	0.5	182	48.7	41.2	7.0	0.5	4.3	59	1.4	5.7	3.8
UK	1,806	2.6	2.1	0.5	0.1	9	2.4	1.6	0.5	0.3	5.0	2	1.1	6.1	2.8
EU15 (excluding Ireland & UK)	506	0.7	0.6	0.1	0.0	1	0.3	0.3	0.0	0.0	2.0	1	2.0	3.9	2.0
EU15 to EU27 Accession States	3,206	4.6	3.7	0.8	0.1	10	2.7	2.7	0.0	0.0	3.1	1	0.3	3.4	3.1
Rest of Europe	508	0.7	0.7	0.0	0.0	4	1.1	0.8	0.3	0.0	7.8	1	2.0	9.8	9.8
Africa	2,060	3.0	2.8	0.2	0.0	12	3.2	3.2	0.0	0.0	5.8	6	2.9	8.7	6.3
Asia	1,971	2.8	2.8	0.1	0.0	13	3.5	3.2	0.3	0.0	6.6	5	2.5	9.1	7.6
America	476	0.7	0.6	0.1	0.0	4	1.1	1.1	0.0	0.0	8.3	1	2.1	10.4	8.4
Australia	105	0.2	0.1	0.0	0.0	1	0.3	0.3	0.0	0.0	9.4	3	28.6	37.7	28.6
New Zealand and Other Oceania	73	0.1	0.1	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Multi-Nationality	1	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Other Nationality	3	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Not Stated	455	0.7	0.7	0.0	0.0	18	4.8	4.5	0.0	0.3	38.1	5	11.0	48.6	28.1
Not Applicable	16,295	23.5	0.0	22.4	1.1	120	32.1	0.0	30.2	1.9	7.3	34	2.1	9.4	7.6
Total	69,318	100	66.2	31.9	1.8	374	100	58.8	38.2	2.9	5.4	118	1.7	7.1	5.1

\* Includes; divorced, married but separated, widowed, and not stated. Notes: See Appendix F for a description of the classification system used for nationality. Father's Nationality

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Father's Nationality	Live Births	Per Cent		t (%) Live B r's Marital		Stillbirths	Per Cent (%)		t (%) Stillbi r's Marital		Stillbirth Rate	Early Neonatal	Early Neonatal	Perinatal Mortality	Adjusted PMR
		(%) Live Births	Married	Never Married	Other*		Stillbirths	Married	Never Married	Other*		Deaths	Mortality Rate	Rate	
Ireland	1,513	67.2	61.5	5.0	0.6	12	63.2	63.2	0.0	0.0	7.9	9	5.9	13.8	11.2
UK	64	2.8	2.6	0.3	0.0	1	5.3	5.3	0.0	0.0	15.4	0	0.0	15.4	15.4
EU15 (excluding Ireland & UK)	32	1.4	1.2	0.2	0.1	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
EU15 to EU27 Accession States	67	3.0	2.3	0.6	0.1	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Rest of Europe	14	0.6	0.5	0.0	0.1	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Africa	59	2.6	2.5	0.0	0.1	1	5.3	5.3	0.0	0.0	16.7	0	0.0	16.7	16.7
Asia	28	1.2	1.2	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
America	14	0.6	0.6	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Australia	6	0.3	0.3	0.0	0.0	0	0.0	0.0	0.0	0.0	0.0	0	0.0	0.0	0.0
Not Stated	31	1.4	1.4	0.0	0.0	1	5.3	5.3	0.0	0.0	31.3	0	0.0	31.3	31.3
Not Applicable	424	18.8	0.0	17.5	1.3	4	21.1	0.0	15.8	5.3	9.3	5	11.8	21.0	21.0
Total	2,252	100	74.1	23.6	2.4	19	100	78.9	15.8	5.3	8.4	14	6.2	14.5	12.8

Notes: \* Includes; divorced, married but separated, widowed, and not stated. See note to Table 3.6a.

### **MATERNAL PARITY**

Maternal parity is a mother's total number of previous live births and stillbirths. Almost 42 per cent of total births in 2007 were to primiparous women<sup>15</sup> and 31.9 per cent were to women who had one previous live birth or stillbirth (see Tables 3.7a and 3.7b).

Over the period 2003 to 2007, the average maternal age rose from 28.2 to 28.9 years for primiparous mothers and from 32.2 to 32.7 years for multiparous mothers. <sup>16</sup> During this five year period the proportion of mothers giving birth for the first time and aged 35 years or more increased from 12 to 14.5 per cent, while the proportion of multiparous mothers aged 35 years or more increased from 31.3 to 35.3 per cent (see Figures 3.8 and 3.9).

Maternal parity by mother's age for 2007 is shown in Figure 3.10. Over 90 per cent of mothers aged less than 20 years gave birth for the first time with only 7.8 per cent delivering for the second time and 0.8 per cent delivering more than a third time. A closer to equal distribution was observed for mothers aged between 30 and 34 years giving birth for the first and second time. As expected, mothers aged 40 years or more were relatively evenly distributed for first, second and third time deliveries.

**TABLE 3.7a: SINGLETON BIRTHS Maternal Parity** Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Maternal Parity	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
0	28,998	41.8	178	6.1	47	1.6	7.7	6.0
1	22,092	31.9	97	4.4	36	1.6	6.0	4.1
2	11,554	16.7	45	3.9	26	2.3	6.1	4.1
3	4,317	6.2	26	6.0	5	1.2	7.1	4.2
4	1,393	2.0	16	11.4	3	2.2	13.5	9.3
5	543	0.8	4	7.3	1	1.8	9.1	7.3
6	220	0.3	5	22.2	0	0.0	22.2	13.5
More than 6	183	0.3	3	16.1	0	0.0	16.1	10.8
Not Stated	18	0.0	0	0.0	0	0.0	0.0	0.0
Total	69,318	100	374	5.4	118	1.7	7.1	5.1

 $<sup>^{15}</sup>$  Women giving birth for the first time


<sup>&</sup>lt;sup>16</sup> Women who have had at least one previous pregnancy resulting in a live birth or stillbirth

**Maternal Parity** 

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007


Maternal Parity	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
0	973	43.2	7	7.1	8	8.2	15.3	14.3
1	742	32.9	7	9.3	5	6.7	16.0	12.1
2	338	15.0	5	14.6	1	3.0	17.5	17.5
3	125	5.6	0	0.0	0	0.0	0.0	0.0
4	46	2.0	0	0.0	0	0.0	0.0	0.0
5	12	0.5	0	0.0	0	0.0	0.0	0.0
6	8	0.4	0	0.0	0	0.0	0.0	0.0
More than 6	8	0.4	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

**FIGURE 3.8**Total Births to Primiparous Mothers in each Maternal Age Group, 2003 and 2007




Notes: See Appendix H, Table H8 for data. 'Not Stated' values are not included in the calculation of percentages.

FIGURE 3.9 Total Births to Multiparous Mothers in each Maternal Age Group, 2003 and 2007



Notes: See Appendix H, Table H9 for data. 'Not Stated' values are not included in the calculation of percentages.

**FIGURE 3.10** Parity by Age of Mother, Total Births, 2007



Notes: See Appendix H, Table H10 for data. 'Not Stated' values are not included in the calculation of percentages. In 2007, of total births to multiparous women, 98.3 per cent recorded no previous stillbirths while 1.7 per cent recorded at least 1 previous stillbirth. This shows very little variation from previous years.

**TABLE 3.8a: SINGLETON BIRTHS** 

Previous Stillbirths (for women having second and subsequent births) Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Previous Stillbirths	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
0	39,636	98.3	187	4.7	69	1.7	6.4	4.3
1	625	1.6	9	14.2	2	3.2	17.4	11.1
2	38	0.1	0	0.0	0	0.0	0.0	0.0
3 and over	12	0.0	0	0.0	0	0.0	0.0	0.0
Not Stated	9	0.0	0	0.0	0	0.0	0.0	0.0
Total	40,320	100	196	4.8	71	1.8	6.6	4.4

Note: Primiparae (i.e. women with no previous live births or stillbirths) are not included in this table.

### **TABLE 3.8b: MULTIPLE BIRTHS**

Previous Stillbirths (for women having second and subsequent births) Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Previous Stillbirths	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
0	1,256	98.2	12	9.5	6	4.8	14.2	11.9
1	19	1.5	0	0.0	0	0.0	0.0	0.0
2	4	0.3	0	0.0	0	0.0	0.0	0.0
Total	1,279	100	12	9.3	6	4.7	13.9	11.6

Note: See note to Table 3.8a.

### **INTERVAL SINCE LAST BIRTH**

Tables 3.9a and 3.9b outline the interval in years since mother's last birth for singleton and multiple births respectively. Just over 42 per cent of total births were first deliveries, a marginal increase from 41.1 per cent in 2003.<sup>17</sup> Of the remaining births, there was a decrease in the proportion of mothers whose previous birth was less than 3 years ago, from 30.5 per cent in 2003 to 29.5 per cent in 2007.

**TABLE 3.9a: SINGLETON BIRTHS** Interval in Years Since Last Birth Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Interval Since Last Birth	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
No previous births	28,998	41.8	178	6.1	47	1.6	7.7	6.0
1 year or less	529	0.8	4	7.5	7	13.2	20.6	11.4
>1 year to 2 years	9,631	13.9	49	5.1	18	1.9	6.9	4.2
>2 years to 3 years	10,142	14.6	35	3.4	14	1.4	4.8	3.5
>3 years to 4 years	6,260	9.0	26	4.1	9	1.4	5.6	3.7
>4 years to 5 years	3,997	5.8	16	4.0	5	1.3	5.2	3.0
>5 years to 6 years	2,704	3.9	15	5.5	4	1.5	7.0	4.4
More than 6 years	6,454	9.3	45	6.9	11	1.7	8.6	6.5
Number of previous births unknown	3	0.0	0	0.0	0	0.0	0.0	0.0
Not Stated	600	0.9	6	9.9	3	5.0	14.9	11.6
Total	69,318	100	374	5.4	118	1.7	7.1	5.1

**Table 3.9b: MULTIPLE BIRTHS** Interval in Years Since Last Birth Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Interval Since Last Birth	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
No previous births	973	43.2	7	7.1	8	8.2	15.3	14.3
1 year or less	30	1.3	0	0.0	0	0.0	0.0	0.0
>1 year to 2 years	308	13.7	2	6.5	0	0.0	6.5	6.5
>2 years to 3 years	302	13.4	2	6.6	2	6.6	13.2	6.6
>3 years to 4 years	210	9.3	4	18.7	1	4.8	23.4	18.8
>4 years to 5 years	92	4.1	0	0.0	0	0.0	0.0	0.0
>5 years to 6 years	61	2.7	0	0.0	2	32.8	32.8	32.8
More than 6 years	266	11.8	4	14.8	1	3.8	18.5	18.5
Not Stated	10	0.4	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

<sup>&</sup>lt;sup>17</sup> For data on 2003, see *Report on Perinatal Statistics for 2003*, The Economic and Social Research Institute, December 2006.

### **BIRTHWEIGHT**

The birthweight of a baby is widely used as a key indicator of health status. According to the World Health Organisation (WHO), low birthweight babies are defined as weighing less than 2,500 grams. This is further divided into very low birthweight (weighing less than 1,500 grams) and extremely low birthweight (weighing less than 1,000 grams). Low birthweight babies are more likely to have poor perinatal outcomes with consequences for later life experience. In 2007, the average birthweight of live births was 3,499 grams for singletons and 2,421 grams for multiples. The average birthweight of total live births in 2007 was 3,465 grams. The percentage of total births classified as low birthweight was 5.3 per cent (3.8 per cent of singletons and 50.4 per cent of multiples). There is considerable variation in low birthweight rates between singleton and multiple births (see Tables 3.10a and 3.10b). The percentage of singleton live births weighing less than 1,500 grams was 0.7 per cent compared to 7.5 per cent for multiple births (see Figure 3.11). The combined percentage of total live births with very low and extremely low birthweights (<1,500g) was just under 1 per cent. In 2007, 92 per cent of total live born babies were in the healthy birthweight category 2,500-4,499 grams (see Figure 3.12).

**TABLE 3.10a: SINGLETON BIRTHS**Birthweight
Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Birthweight (grams)	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
500-749	86	0.1	67	437.9	35	407.0	666.7	645.8
750-999	91	0.1	26	222.2	12	131.9	324.8	261.7
1000-1249	131	0.2	30	186.3	5	38.2	217.4	160.0
1250-1499	161	0.2	26	139.0	9	55.9	187.2	105.9
1500-1999	516	0.7	41	73.6	16	31.0	102.3	58.4
2000-2499	1,645	2.4	47	27.8	18	10.9	38.4	19.9
2500-2999	7,462	10.8	45	6.0	10	1.3	7.3	5.5
3000-3499	23,127	33.4	50	2.2	7	0.3	2.5	2.1
3500-3999	24,462	35.3	28	1.1	4	0.2	1.3	1.0
4000-4499	9,658	13.9	10	1.0	2	0.2	1.2	0.9
4500 and Over	1,969	2.8	2	1.0	0	0.0	1.0	1.0
Not Stated	10	0.0	2	166.7	0	0.0	166.7	90.9
Total	69,318	100	374	5.4	118	1.7	7.1	5.1

<sup>&</sup>lt;sup>18</sup> EURO-PERISTAT Project, with SCPE, EUROCAT, EURONEOSTAT. European Perinatal Health Report, 2008. www.europeristat.com


**TABLE 3.10b: MULTIPLE BIRTHS** 

Birthweight

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007


Birthweight (grams)	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
500-749	19	0.8	1	50.0	7	368.4	400.0	400.0
750-999	41	1.8	5	108.7	5	122.0	217.4	200.0
1000-1249	48	2.1	0	0.0	1	20.8	20.8	0.0
1250-1499	62	2.8	0	0.0	0	0.0	0.0	0.0
1500-1999	301	13.4	2	6.6	0	0.0	6.6	6.6
2000-2499	665	29.5	1	1.5	0	0.0	1.5	1.5
2500-2999	766	34.0	4	5.2	0	0.0	5.2	5.2
3000-3499	304	13.5	2	6.5	0	0.0	6.5	6.5
3500-3999	44	2.0	0	0.0	1	22.7	22.7	0.0
4000-4499	2	0.1	0	0.0	0	0.0	0.0	0.0
Not Stated	0	0.0	4	1000.0	0	0.0	1,000.0	1,000.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

**FIGURE 3.11** Birthweights: Cumulative Percentages of Total Live Births, 2007



Note: See Tables 3.10a and 3.10b for data. 'Not Stated' values are not included in the calculation of percentages.


**FIGURE 3.12** Percentage Births by Birthweight Groups, Total Live Births, 2007



Note: See Appendix H, Table H11 for data. 'Not Stated' values are not included in the calculation of percentages.

The average birthweight of stillborn babies was 2,012 grams in 2007 compared to the 3,465 grams reported for live born babies. Low birthweight occurred in 63.6 per cent of stillborn babies. Just over 40 per cent of stillborn babies weighed less than 1,500 grams. Figure 3.13 shows the percentage distribution across birthweight groups by sex. A marginally smaller proportion of male live born babies were low birthweight (4.8 per cent) compared with female babies (5.7 per cent). The average birthweight of live born male babies was 3,529 grams compared to 3,399 grams for females.

**FIGURE 3.13** Percentage Distribution for Males and Females across Birthweight Groups, Total Births, 2007



Note: See Appendix H, Table H12 for data. 'Not Stated' values are not included in the calculation of percentages.

### **GESTATIONAL AGE**

The average gestational age for total births in 2007 was 39.3 weeks. The percentage of all babies born at term (37-41 weeks) was 90.5 per cent (see Tables 3.11a and 3.11b). Preterm births (born before 37 completed weeks gestation) were associated with higher risk of perinatal death and comprise 6.1 per cent of total births. Figure 3.14 presents perinatal mortality rates by gestational age for total births. Most preterm births occurred at a gestational age of 32-36 completed weeks (4.9 per cent of total births). For stillborn babies the mean gestational age was 33.3 weeks compared to 39.4 weeks for live borns. Preterm birth occurred in 59.8 per cent of stillborns and 5.8 per cent of live borns. For singleton births the average gestational age was 39.4 completed weeks compared with 35.8 for twins and 32.5 for higher order multiple births. Preterm delivery occurred in 46.3 per cent of twins and in 96.5 per cent of higher order multiple births. Mortality rates should be interpreted with caution as they can differ significantly for singleton and multiple births.


**TABLE 3.11a: SINGLETON BIRTHS** Gestational Age at Delivery Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Gestational Age (weeks)	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Less than 22	1	0.0	5	833.3	1	1,000.0	1,000.0	1,000.0
22-27	163	0.2	70	300.4	41	251.5	476.4	452.9
28-31	368	0.5	64	148.1	16	43.5	185.2	139.4
32-36	2,535	3.7	84	32.1	40	15.8	47.3	24.2
37-41	63,784	92.0	148	2.3	20	0.3	2.6	2.0
42 and Over	2,457	3.5	3	1.2	0	0.0	1.2	0.8
Not Stated	10	0.0	0	0.0	0	0.0	0.0	0.0
Total	69,318	100	374	5.4	118	1.7	7.1	5.1

**TABLE 3.11b: MULTIPLE BIRTHS** Gestational Age at Delivery Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Gestational Age (weeks)	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Less than 22	0	0.0	2	1000.0	0	0.0	1,000.0	1,000.0
22-27	49	2.2	2	39.2	12	244.9	274.5	260.0
28-31	147	6.5	1	6.8	0	0.0	6.8	6.8
32-36	887	39.4	7	7.8	1	1.1	8.9	7.8
37-41	1,167	51.8	7	6.0	1	0.9	6.8	5.1
42 and Over	2	0.1	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

**FIGURE 3.14** Perinatal Mortality Rates by Gestational Age at Delivery, Total Births, 2007



Notes: See Appendix H, Table H13 for data. Rates based on 20 or less observations are not included in this Figure.

### **MONTH OF BIRTH**

There was some seasonal variation in the distribution of singleton births by month. As shown in Table 3.12a fewer babies were born during the winter months (January and February) and higher numbers were born during July to October, peaking in August (9.1 per cent) and September (8.9 per cent). Multiple births show higher percentages born during the later part of the year (see Table 3.12b and Figure 3.15).

**TABLE 3.12a: SINGLETON BIRTHS** Month of Birth Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Month of Birth	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
January	5,430	7.8	29	5.3	12	2.2	7.5	4.4
February	4,848	7.0	25	5.1	8	1.7	6.8	4.5
March	5,657	8.2	25	4.4	10	1.8	6.2	4.4
April	5,476	7.9	27	4.9	6	1.1	6.0	4.7
May	5,975	8.6	28	4.7	14	2.3	7.0	5.2
June	5,813	8.4	34	5.8	8	1.4	7.2	4.6
July	6,030	8.7	33	5.4	10	1.7	7.1	5.3
August	6,285	9.1	34	5.4	12	1.9	7.3	5.2
September	6,173	8.9	38	6.1	4	0.6	6.8	5.5
October	6,030	8.7	28	4.6	9	1.5	6.1	4.1
November	5,743	8.3	39	6.7	14	2.4	9.2	6.8
December	5,858	8.5	34	5.8	11	1.9	7.6	6.1
Total	69,318	100	374	5.4	118	1.7	7.1	5.1


**TABLE 3.12b: MULTIPLE BIRTHS** 

Month of Birth

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Month of Birth	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
January	182	8.1	3	16.2	0	0.0	16.2	16.2
February	134	6.0	1	7.4	1	7.5	14.8	14.8
March	163	7.2	1	6.1	2	12.3	18.3	6.2
April	139	6.2	0	0.0	1	7.2	7.2	7.2
May	200	8.9	1	5.0	0	0.0	5.0	5.0
June	164	7.3	2	12.0	0	0.0	12.0	12.0
July	217	9.6	1	4.6	0	0.0	4.6	4.6
August	242	10.7	3	12.2	4	16.5	28.6	28.6
September	198	8.8	1	5.0	1	5.1	10.1	10.1
October	206	9.1	1	4.8	4	19.4	24.2	19.4
November	229	10.2	1	4.3	1	4.4	8.7	8.7
December	178	7.9	4	22.0	0	0.0	22.0	16.6
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

**FIGURE 3.15** Percentage of Births by Month of Birth, Total Births, 2007



Note: See Tables 3.13a and 3.13b for data.

# **SECTION FOUR**

# **PERINATAL CARE**



### **INTRODUCTION**

This section presents a detailed review on aspects of perinatal care for the mother and baby. Singleton and multiple births are reported in separate tables. The following tables outline data for live births, stillbirths and early neonatal deaths, together with the relevant rates. The perinatal mortality rate and adjusted perinatal mortality rate are also outlined in each table. Tables and figures are presented in some detail for a number of variables, including type of antenatal care, timing of first antenatal visit, mother's antenatal and postnatal length of stay, method of delivery, mother's immunity to rubella, infant's length of stay, and type of feeding.

### **TYPE OF ANTENATAL CARE**

The preferred choice of antenatal care was the combined hospital and G.P. option accounting for just over 77 per cent of total births. 19 Hospital/Obstetrician care accounted for a further 20.8 per cent of total births, leaving only a very small proportion of births delivered under other antenatal care schemes. Tables 4.1a and 4.1b show that a higher percentage of multiple births are delivered under hospital/obstetrician schemes when compared to singleton births. Deliveries under domino schemes and midwifery led maternity units within some hospitals were included under combined care.

**TABLE 4.1a: SINGLETON BIRTHS** Type of Antenatal Care Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Type of Antenatal Care	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Hospital/Obstetrician	14,059	20.3	117	8.3	30	2.1	10.4	6.8
General Practitioner Only	695	1.0	5	7.1	3	4.3	11.4	8.6
Hospital and G.P. Combined	53,835	77.7	238	4.4	80	1.5	5.9	4.3
None	179	0.3	10	52.9	4	22.3	74.1	69.1
Midwife Only	503	0.7	3	5.9	1	2.0	7.9	7.9
Unknown	47	0.1	1	20.8	0	0.0	20.8	20.8
Total	69,318	100	374	5.4	118	1.7	7.1	5.1

**TABLE 4.1b: MULTIPLE BIRTHS** Type of Antenatal Care Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Type of Antenatal Care	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Hospital/Obstetrician	755	33.5	9	11.8	6	7.9	19.6	17.1
General Practitioner Only	18	0.8	0	0.0	0	0.0	0.0	0.0
Hospital and G.P. Combined	1,474	65.5	10	6.7	8	5.4	12.1	10.8
None	4	0.2	0	0.0	0	0.0	0.0	0.0
Unknown	1	0.0	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

<sup>&</sup>lt;sup>19</sup> Throughout this section, 'Not Stated' values are not included in the calculation of percentages for discussion.

### **DURATION OF PREGNANCY AT FIRST ANTENATAL VISIT**

Although 77.2 per cent of total births were recorded as receiving combined antenatal care in 2007, the date of the first visit to the doctor was recorded as 'not known' for 45.1 per cent of these births. As a result of the absence of these data, the timing of first contact with health professionals within this category will reflect the date of the first hospital visit, even though this is likely to have been later than the first doctor visit. The percentage of mothers attending antenatal care during the first 12 weeks in 2007 was, therefore, estimated at around 55 per cent of total births, which is low by European standards. Consequently, caution should be exercised in interpreting the data presented here on the timing of the first visit to health services (see Tables 4.2a and 4.2b).

**TABLE 4.2a: SINGLETON BIRTHS** Time of First Visit to Doctor or Hospital during Pregnancy Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Time of first visit (weeks)	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
0-12	36,633	52.8	162	4.4	61	1.7	6.1	4.1
13-19	22,167	32.0	130	5.8	31	1.4	7.2	5.3
20 and over	7,910	11.4	66	8.3	18	2.3	10.5	8.0
Not Stated	2,608	3.8	16	6.1	8	3.1	9.1	7.6
Total	69,318	100	374	5.4	118	1.7	7.1	5.1

**TABLE 4.2b: MULTIPLE BIRTHS** Time of First Visit to Doctor or Hospital during Pregnancy Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Time of first visit (weeks)	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
0-12	1,308	58.1	5	3.8	5	3.8	7.6	6.1
13-19	636	28.2	11	17.0	6	9.4	26.3	24.8
20 and over	206	9.1	0	0.0	3	14.6	14.6	9.8
Not Stated	102	4.5	3	28.6	0	0.0	28.6	28.6
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

### **MOTHER'S ANTENATAL LENGTH OF STAY**

The average antenatal length of stay was 0.7 days for singleton births and 2.1 days for multiple births. Tables 4.3a and 4.3b outline the distribution of antenatal length of stay for singleton and multiple births. Over 90 per cent of singleton births and 77.3 per cent of multiple births were delivered within less than 2 days of mother's admission to hospital. Mother's antenatal length of stay was 9 days or more for only 0.8 per cent of singleton births and 5.8 per cent for multiple births.

**TABLE 4.3a: SINGLETON BIRTHS** Mother's Antenatal Length of Stay Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Antenatal Length of Stay	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
0-1 Days	62,349	90.2	294	4.7	97	1.6	6.2	4.5
2 Days	3,735	5.4	43	11.4	10	2.7	14.0	8.0
3-5 Days	1,977	2.9	26	13.0	4	2.0	15.0	13.5
6-8 Days	477	0.7	6	12.4	3	6.3	18.6	14.6
9-11 Days	186	0.3	1	5.3	0	0.0	5.3	5.3
12-14 Days	106	0.2	0	0.0	0	0.0	0.0	0.0
More than 2 Weeks	251	0.4	3	11.8	4	15.9	27.6	23.7
Not Stated	52	0.1	0	0.0	0	0.0	0.0	0.0
Total	69,133	100	373	5.4	118	1.7	7.1	5.1

'Not Stated' includes 41 births where place of birth was recorded as domiciliary but birth was registered by a Notes: hospital. For these cases, the mother was not admitted, or was admitted post delivery, under a planned community midwife scheme. Therefore, dates of mother's antenatal length of stay were not applicable. There were 5 births where place of birth was recorded as 'Born Before Arrival'. In these cases, mother's admission date was after the infant's date of birth. Therefore, antenatal length of stay was not applicable. There were 6 births where mother's

Domiciliary births, of which there were 186 singleton births, are excluded from this table. See Appendix G for detailed analysis.

**TABLE 4.3b: MULTIPLE BIRTHS** Mother's Antenatal Length of Stay Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Antenatal Length of Stay	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
0-1 Days	1,740	77.3	16	9.1	9	5.2	14.2	12.5
2 Days	121	5.4	0	0.0	0	0.0	0.0	0.0
3-5 Days	181	8.0	3	16.3	3	16.6	32.6	32.6
6-8 Days	78	3.5	0	0.0	0	0.0	0.0	0.0
9-11 Days	27	1.2	0	0.0	0	0.0	0.0	0.0
12-14 Days	26	1.2	0	0.0	1	38.5	38.5	38.5
More than 2 weeks	79	3.5	0	0.0	1	12.7	12.7	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

Note: There were no domiciliary multiple births.

### **MOTHER'S POSTNATAL LENGTH OF STAY**

Maternal postnatal length of stay can be influenced by factors such as method of delivery, post delivery complications, neonatal mortality, and hospital discharge policies. In 2007, the mother's median postnatal length of stay was 3 days for live born singleton births and 4 days for live born multiple births. Average postnatal lengths of stay for singleton and multiple births were 2.8 and 4.6 days respectively.

For singleton births, 48 per cent of mothers remained in hospital for less than 3 days and a further 48.8 per cent were discharged between 3 and 5 days (see Table 4.4a). For multiple births, 8.7 per cent of mothers remained in hospital in hospital for less than 3 days, whilst 72.2 per cent were discharged between 3 and 5 days post delivery (see Table 4.4b).

**TABLE 4.4a: SINGLETON BIRTHS** Mother's Postnatal Length of Stay Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Postnatal Length of Stay	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
0-1 Days	12,097	17.5	242	19.6	54	4.5	24.0	17.9
2 Days	20,968	30.3	67	3.2	19	0.9	4.1	3.0
3-5 Days	33,837	48.9	53	1.6	42	1.2	2.8	1.8
6-8 Days	1,910	2.8	4	2.1	2	1.0	3.1	1.6
9-11 Days	204	0.3	4	19.2	0	0.0	19.2	14.5
12-14 Days	33	0.0	0	0.0	0	0.0	0.0	0.0
More than 2 Weeks	42	0.1	2	45.5	1	23.8	68.2	23.8
Not Stated	42	0.1	1	23.3	0	0.0	23.3	23.3
Total	69,133	100	373	5.4	118	1.7	7.1	5.1

Notes:

'Not Stated' includes 37 births where place of birth was recorded as domiciliary but birth was registered by a hospital. For these cases, the mother was not admitted under a planned community midwife scheme. Therefore, dates of mother's postnatal length of stay were not applicable. There were 6 births where mother's discharge date

Domiciliary births, of which there were 186 singleton births, are excluded from this table. See Appendix G for detailed analysis.


**TABLE 4.4b: MULTIPLE BIRTHS** Mother's Postnatal Length of Stay Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Postnatal Length of Stay	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
0-1 Days	33	1.5	0	0.0	5	151.5	151.5	125.0
2 Days	163	7.2	1	6.1	3	18.4	24.4	24.4
3-5 Days	1,621	72.0	16	9.8	4	2.5	12.2	10.4
6-8 Days	370	16.4	0	0.0	2	5.4	5.4	5.4
9-11 Days	43	1.9	2	44.4	0	0.0	44.4	44.4
12-14 Days	17	0.8	0	0.0	0	0.0	0.0	0.0
More than 2 weeks	5	0.2	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

Note: There were no domiciliary multiple births.


The average postnatal length of stay for mothers differed by method of delivery and plurality. Average maternal postnatal length of stay for spontaneous delivery for singleton births was 2.1 days compared to 3.4 days for multiple births. Singleton births delivered by caesarean section resulted in an average maternal postnatal length of stay of 4.4 days compared to 5.2 days for multiple births. Figures 4.1 and 4.2 show the postnatal lengths of stay by method of delivery for singleton and multiple births. For spontaneous deliveries, 68.5 per cent of singleton births compared to 23.7 per cent of multiple births recorded a postnatal length of stay of less than 3 days, and a length of stay between 3 and 5 days for 30.6 per cent for singletons and 68.7 per cent for multiple births. Delivery by caesarean section with a postnatal length of stay of 6 days or more accounted for 9.1 per cent of singleton caesarean deliveries compared to 24.7 per cent for multiple caesarean deliveries.

**FIGURE 4.1** Mother's Postnatal Length of Stay by Method of Delivery, Singleton Births, 2007



See Appendix H, Table H14 for data. 'Not Stated' values are not included in the calculation of percentages. Note:

FIGURE 4.2 Mother's Postnatal Length of Stay by Method of Delivery, Multiple Births, 2007



Note: See Appendix H, Table H15 for data. 'Not Stated' values are not included in the calculation of percentages.

### **ADVANCE HOSPITAL BOOKINGS**

In 2007, 99 per cent of total births were booked at time of admission to hospital. A smaller proportion of multiple births, 97.3 per cent, were booked at admission compared to singleton births, 99.1 per cent (see Tables 4.5a and 4.5b).

**TABLE 4.5a: SINGLETON BIRTHS** 

**Advance Hospital Bookings** 

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Advance Hospital Bookings	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Booked	68,474	99.0	350	5.1	108	1.6	6.7	4.7
Not Booked	601	0.9	22	35.3	10	16.6	51.4	43.7
Not Stated	58	0.1	1	16.9	0	0.0	16.9	16.9
Total	69,133	100	373	5.4	118	1.7	7.1	5.1

Notes:

'Not Stated' includes 36 births where place of birth was recorded as domiciliary but birth was registered by a hospital. For these cases, the mother was not admitted under a planned community midwife scheme. Therefore, hospital bookings were not applicable.

Domiciliary births, of which there were 186 singleton births, are excluded from this table. See Appendix G for detailed analysis.

### **TABLE 4.5b MULTIPLE BIRTHS**

**Advance Hospital Bookings** 

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Advance Hospital Bookings	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Booked	2,190	97.2	19	8.6	14	6.4	14.9	13.2
Not Booked	62	2.8	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

Note: There were no domiciliary multiple births.

#### **RUBELLA: IMMUNE STATUS OF MOTHER**

In 2007, of total babies born, 92.4 per cent of births were to mothers with rubella immunity. A further 2.9 per cent of births were to mothers who were not rubella immune and mother's rubella immunity was not known for 4.8 per cent of total babies born (see Tables 4.6a and 4.6b).

**TABLE 4.6a: SINGLETON BIRTHS** Rubella: Immune Status of Mother, Numbers and Percentages, 2007

Rubella	Number (Live Births and Stillbirths)	Per Cent (%)
Immune	64,372	92.4
Not Immune	2,024	2.9
Unknown	3,294	4.7
Not Stated	2	0.0
Total	69,692	100

TABLE 4.6b: MULTIPLE BIRTHS
Rubella: Immune Status of Mother,
Numbers and Percentages, 2007

Rubella	Number (Live Births and Stillbirths)	Per Cent (%)
Immune	2,092	92.1
Not Immune	47	2.1
Unknown	132	5.8
Total	2,271	100

### **METHOD OF DELIVERY**

Among babies born in Ireland, the percentage of normal vaginal deliveries for live born babies decreased from 61.2 per cent in 2003 to 57.3 per cent in 2007. Tables 4.7a and 4.7b outline method of delivery for singleton and multiple births. There were 18,781 live births delivered by caesarean section, representing a caesarean section rate of 26.2 per cent for total live births in 2007, 25 per cent for singletons and 63.7 per cent for multiples, (see Figure 4.3). In 2007, 15.8 per cent of total live births were born by instrumental delivery (either forceps or vacuum extraction). These percentages differed for live singleton births (16 per cent) and live multiple births (9.5 per cent).

**TABLE 4.7a: SINGLETON BIRTH** 

Method of Delivery

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Method of Delivery	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Spontaneous	40,484	58.4	297	7.3	61	1.5	8.8	6.5
Breech / Forceps	259	0.4	37	125.0	8	30.9	152.0	109.9
Forceps	2,588	3.7	6	2.3	2	0.8	3.1	2.7
Vacuum extraction	8,523	12.3	6	0.7	1	0.1	0.8	0.8
Caesarean section	17,348	25.0	28	1.6	45	2.6	4.2	2.5
Other specified & combined	111	0.2	0	0.0	1	9.0	9.0	0.0
Not stated	5	0.0	0	0.0	0	0.0	0.0	0.0
Total	69,318	100	374	5.4	118	1.7	7.1	5.1


# **TABLE 4.7b: MULTIPLE BIRTHS**

Method of Delivery

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Method of Delivery	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Spontaneous	492	21.8	6	12.0	4	8.1	20.1	20.1
Breech / Forceps	102	4.5	2	19.2	2	19.6	38.5	38.5
Forceps	62	2.8	0	0.0	2	32.3	32.3	32.3
Vacuum extraction	151	6.7	0	0.0	1	6.6	6.6	6.6
Caesarean section	1,433	63.6	11	7.6	5	3.5	11.1	8.3
Other specified & combined	11	0.5	0	0.0	0	0.0	0.0	0.0
Not stated	1	0.0	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	19	8.4	14	6.2	14.5	12.8

FIGURE 4.3 Percentage of Births by Method of Delivery for Live Singleton and Live Multiple Births, 2007



Note: See Tables 4.13 and 4.14 for data. 'Not Stated' values are not included in the calculation of percentages.

The percentage of births born by caesarean section increased with maternal age (see Table 4.8). In 2007, caesarean section rates ranged from 15.7 per cent for mothers aged less than 20 years to 38 per cent for mothers aged 40 years and older. The percentage of babies delivered by spontaneous delivery fell from 64.8 per cent for mothers aged less than 20 years to 50.5 per cent to mothers aged 40 years and older. The percentage delivered by vacuum extraction also fell from 14.8 per cent for mothers aged less than 20 years to 7.5 per cent for mothers aged 40 years and older.

**TABLE 4.8: TOTAL BIRTHS** Method of Delivery by Mother's Age, 2007

Method of Delivery				Mother	's Age			
Frequency	Under	20-24	25-29	30-34	35-39	40 Years	Not	Total
Col. Per Cent	20 Years	Years	Years	Years	Years	and Over	Stated	
Spontaneous	1,634	5,764	10,088	13,507	8,689	1,582	15	41,279
	64.8	64.1	60.4	54.9	54.3	50.5	68.2	57.4
Breech / Forceps	16	50	86	131	97	19	1	400
	0.6	0.6	0.5	0.5	0.6	0.6	4.5	0.6
Forceps	100	329	651	1,011	470	94	1	2,656
	4.0	3.7	3.9	4.1	2.9	3.0	4.5	3.7
Vacuum extraction	373	1,222	2,236	3,064	1,547	236	2	8,680
	14.8	13.6	13.4	12.5	9.7	7.5	9.1	12.1
Caesarean section	397	1,620	3,619	6,835	5,156	1,190	3	18,820
	15.7	18.0	21.7	27.8	32.2	38.0	13.6	26.2
Other specified & combined	1	7	19	49	34	12	0	122
	0.0	0.1	0.1	0.2	0.2	0.4	0.0	0.2
Not stated	0	2	1	1	2	0	0	6
	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Total	2,521	8,994	16,700	24,598	15,995	3,133	22	71,963
	100	100	100	100	100	100	100	100

#### **SIZE OF MATERNITY UNIT**


Over 57 per cent of total births in Ireland take place in maternity units accommodating at least 4,000 births annually. A further 20.6 per cent of total births occur in maternity units reporting between 2,000 and just under 4,000 births annually (see Table 4.9 and Figure 4.4).

**TABLE 4.9: TOTAL BIRTHS**Size of Maternity Unit (number of live births and stillbirths annually)
Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Size of Maternity Unit	Number of Hospitals	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate		Early Neonatal Mortality Rate	Mortality	Adjusted PMR
Less than 999	4	1,991	2.8	10	5.0	3	1.5	6.5	5.0
1000-1999	8	13,438	18.8	68	5.0	24	1.8	6.8	4.5
2000-2999	5	11,304	15.8	64	5.4	18	1.4	6.8	5.1
3000-3999	1	3,401	4.8	16	5.4	3	1.4	6.8	5.1
4000 and Over	6	41,251	57.8	234	5.6	84	2.0	7.7	5.7
Total	24	71,385	100.0	392	5.5	132	1.8	7.3	5.3

Notes: This table includes singleton and multiple births but excludes domiciliary births, of which there were 186 singleton births. There are 20 maternity units in Ireland at present, all of which deliver over 1,000 births per annum. In March 2007, the separate maternity units in Cork transferred their services to the Cork University Maternity Hospital. Occasionally, births can take place in hospitals without maternity units; these are usually exceptional circumstances and are included in the NPRS data set.

**FIGURE 4.4**Percentage of Births by Size of Maternity Unit, Total Live Births, 2007



Note: See Table 4.9 for data. The size of the unit refers to the number of live births and stillbirths occurring in 2007.

#### **INTER-HOSPITAL TRANSFER OF INFANTS**

For singleton births, almost all live births remained in the hospital in which they were born with just less than 1 per cent of live births being transferred to another hospital for medical reasons (see Tables 4.10a). For multiple births, 97 per cent remained in the hospital of birth with almost 3 per cent being transferred for medical reasons (see Table 4.10b).

**TABLE 4.10a: SINGLETON BIRTHS** 

Inter-Hospital Transfer of Infants

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Infant Transfer	Live Births	Per Cent (%) Live Births	Early Neonatal Deaths	Early Neonatal Mortality Rate
Transferred	393	0.6	5	12.7
Not Transferred	68,688	99.4	113	1.6
Not Stated	52	0.1	0	0.0
Total	69,133	100	118	1.7

Notes:

'Not Stated' includes 35 births where place of birth was recorded as domiciliary but birth was registered by a hospital. For these cases, the baby was not admitted under a planned community midwife scheme. Therefore, transfer details were not applicable.

Domiciliary births, of which there were 185 singleton live births, are excluded from this table. See Appendix G for detailed analysis.

**TABLE 4.10b: MULTIPLE BIRTHS** 

Inter-Hospital Transfer of Infants

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Infant Transfer	Live Births	Per Cent (%) Live Births	Early Neonatal Deaths	Early Neonatal Mortality Rate
Transferred	65	2.9	0	0.0
Not Transferred	2,184	97.0	14	6.4
Not Stated	3	0.1	0	0.0
Total	2,252	100	14	6.2

Note: There were no domiciliary multiple births.

#### **INFANT'S LENGTH OF STAY**

Almost 95 per cent of live singleton births and 59 per cent of live multiple births were discharged from hospital within 5 days of birth (see Tables 4.11a and 4.11b respectively). Approximately 50 per cent of both singleton and multiple births were discharged between 3 and 5 days following delivery. For multiple births, a further 16.7 per cent remained in hospital for between 6 and 10 days and a further 17.3 per cent were discharged between 11 and 28 days.

**TABLE 4.11a: SINGLETON BIRTHS** Percentage Distribution of Live Births by Infant's Length of Stay in Hospital, 2007

**TABLE 4.11b: MULTIPLE BIRTHS** Percentage Distribution of Live Births by Infant's Length of Stay in Hospital, 2007


Infant's Length of Stay (Days)	Live Births	Per Cent (%) Live Births
0-2	32,566	47.2
3-5	32,693	47.4
6-10	2,403	3.5
11-28	884	1.3
29 or more	424	0.6
Not Stated	45	0.1
Total	69,015	100

Note:

Infant's Length of Stay (Days)	Live Births	Per Cent (%) Live Births
0-2	148	6.6
3-5	1,172	52.4
6-10	373	16.7
11-28	388	17.3
29 or more	156	7.0
Not Stated	1	0.0
Total	2,238	100

Notes: These tables do not include domiciliary births and early neonatal deaths. 'Not Stated' includes 37 births where place of birth was recorded as domiciliary but birth was registered by a hospital. In these cases, the infant was not admitted under a planned community midwife scheme. Therefore, infant's length of stay was not applicable.

FIGURE 4.5 Percentage of Births by Infant's Length of Stay for Live Singleton and Live Multiple Births, 2007



See Tables 4.11a and 4.11b for data. 'Not Stated' values are not included in the calculation of percentages.

#### **INFANT'S TYPE OF FEEDING**

Breastfeeding statistics are based on live births only and exclude early neonatal deaths. Type of feeding is recorded at the time of discharge from hospital. As shown in Tables 4.12a and 4.12b, artificial feeding is more common for both singleton (49.2 per cent) and multiple births (57.6 per cent). The overall percentage of total live births exclusively breastfed at the time of discharge from hospital in 2007 was 45 per cent. This is slightly different to the percentage of mother's breastfeeding (45.2 per cent), due to plurality of births (see Table 2.1, page 9).

**TABLE 4.12a: SINGLETON BIRTHS** Age of Mother by Infant's Type of Feeding, Numbers and Percentages, Live Births, 2007

Age of Mother			Type of Feeding		
Frequency	Artificial	Breast	Combined	Not Stated	Total
Row Per Cent					
Col. Per Cent					
Under 15 Years	6	4	0	0	10
	60.0	40.0	0.0	0.0	100
	0.0	0.0	0.0	0.0	0.0
15-19 Years	1,776	560	86	6	2,428
	73.1	23.1	3.5	0.2	100
	5.2	1.8	2.3	6.2	3.5
20-24 Years	5,122	3,251	424	17	8,814
	58.1	36.9	4.8	0.2	100
	15.0	10.3	11.5	17.5	12.7
25-29 Years	8,138	7,177	925	18	16,258
	50.1	44.1	5.7	0.1	100
	23.9	22.8	25.1	18.6	23.5
30-34 Years	10,885	11,449	1,286	32	23,652
	46.0	48.4	5.4	0.1	100
	31.9	36.4	34.9	33.0	34.1
35-39 Years	6,856	7,505	798	20	15,179
	45.2	49.4	5.3	0.1	100
	20.1	23.9	21.6	20.6	21.9
40-44 Years	1,229	1,462	164	3	2,858
	43.0	51.2	5.7	0.1	100
	3.6	4.6	4.4	3.1	4.1
45 Years and	46	51	5	1	103
Over	44.7	49.5	4.9	1.0	100
	0.1	0.2	0.1	1.0	0.1
Not Stated	12	4	0	0	16
	75.0	25.0	0.0	0.0	100
	0.0	0.0	0.0	0.0	0.0
Total	34,070	31,463	3,688	97	69,318
	49.2	45.4	5.3	0.1	100
	100	100	100	100	100

Note: The 'Not Stated' category for type of feeding includes 85 early neonatal deaths.

Age of Mother by Infant's Type of Feeding, Numbers and Percentages, Live Births, 2007

Age of Mother			Type of Feeding		
Frequency	Artificial	Breast	Combined	Not Stated	Total
Row Per Cent					
Col. Per Cent					
15-19 Years	52	11	5	1	69
	75.4	15.9	7.2	1.4	100
	4.0	1.6	1.9	12.5	3.1
20-24 Years	74	29	19	3	125
	59.2	23.2	15.2	2.4	100
	5.7	4.2	7.2	37.5	5.6
25-29 Years	196	118	45	0	359
	54.6	32.9	12.5	0.0	100
	15.2	17.1	17.1	0.0	15.9
30-34 Years	506	237	95	3	841
	60.2	28.2	11.3	0.4	100
	39.2	34.4	36.1	37.5	37.3
35-39 Years	382	241	83	1	707
	54.0	34.1	11.7	0.1	100
	29.6	35.0	31.6	12.5	31.4
40-44 Years	78	46	16	0	140
	55.7	32.9	11.4	0.0	100
	6.0	6.7	6.1	0.0	6.2
45 Years and	4	7	0	0	11
Over	36.4	63.6	0.0	0.0	100
	0.3	1.0	0.0	0.0	0.5
Total	1,292	689	263	8	2,252
	57.4	30.6	11.7	0.4	100
	100	100	100	100	100


Note: The 'Not Stated' category for type of feeding includes 7 early neonatal deaths.

As shown in Figure 4.6, the percentage of total live births breastfed varies by maternal age. Infants born to older mothers are more likely to be breastfed than those born to younger mothers. In 2007, 23 per cent of infants born to mothers under 20 were exclusively breastfeeding compared to 50.4 per cent born to those mothers aged 40-44 and 51.3 per cent to those aged 45 and over.

Figure 4.7 shows that the prevalence of breastfeeding at time of discharge also varies by geographic location. Infants born to mothers resident in Dublin County recorded the highest rate of breastfeeding at 54.8 per cent, closely followed by Westmeath with 52.3 per cent. Very low rates were recorded for infants whose mother was resident in Limerick and Clare (21.6 and 23.4 per cent respectively).


Note:

FIGURE 4.6 Percentage of Breastfed Infants by Age of Mother, Total Live Births, 2007



Note: See Appendix H, Table H16 for data. 'Not Stated' values are not included in the calculation of percentages.

FIGURE 4.7 Percentage Distribution of Breastfed Infants by Mother's County of Residence, Total Live Births, 2007



See Appendix H, Table H17 for data. 'Not Stated' values are not included in the calculation of percentages. Example: Of all live births to mothers resident in Galway, 41.2 per cent were breastfed.


Type of feeding was also found to vary according to mother's occupation (see Table 4.13 and Figure 4.8). Within all occupational categories combined feeding accounted for less than 10 per cent of total live births. In general, infants born to mothers in the higher occupational groupings were more likely to be breastfed while infants born to mothers in the lower occupational groupings were more likely to be fed artificially. In 2007, 64.2 per cent of infants born to mothers who were unemployed were fed artificially, while 63.8 per cent of infants born to mothers with a higher professional occupation were breastfed.

**TABLE 4.13: TOTAL LIVE BIRTHS** Mother's Occupation by Infant's Type of Feeding, 2007

Mother's Occupation			Type of Feeding		
Frequency	Artificial	Breast	Combined	Not Stated	Total
Row Per Cent					
Col. Per Cent					
Farmers & Farm Managers	29	34	2	0	65
	44.6	52.3	3.1	0.0	100
	0.1	0.1	0.1	0.0	0.1
Other Agricultural	36	50	7	0	93
Occupations & Fishermen	38.7	53.8	7.5	0.0	100
Higher Professional	0.1 1,316	0.2 2,754	0.2 248	0.0 5	0.1 4,323
nigher Professional	30.4	63.7	5.7	0.1	100
	3.7	8.6	6.3	4.8	6.0
Lower Professional	3,198	6,101	591	15	9,905
20Wei Froressional	32.3	61.6	6.0	0.2	100
	9.0	19.0	15.0	14.3	13.8
Employers & Managers	2,186	2,648	280	7	5,121
	42.7	51.7	5.5	0.1	100
	6.2	8.2	7.1	6.7	7.2
Salaried Employees	509	515	58	3	1,085
	46.9	47.5	5.3	0.3	100
	1.4	1.6	1.5	2.9	1.5
Intermediate Non-Manual	9,193	6,455	702	21	16,371
Workers	56.2	39.4	4.3	0.1	100
	26.0	20.1	17.8	20.0	22.9
Other Non-Manual Workers	4,780	3,631	483	10	8,904
	53.7	40.8	5.4	0.1	100
Chilled Manual Markers	13.5	11.3	12.2	9.5	12.4
Skilled Manual Workers	453	742	86	2	1,283
	35.3 1.3	57.8 2.3	6.7 2.2	0.2 1.9	100 1.8
Semi-Skilled Manual Workers	1,340	978	126	1.9	2,448
Seriii-Skiiied Waridar Workers	54.7	40.0	5.1	0.2	100
	3.8	3.0	3.2	3.8	3.4
Unskilled Manual Workers	1	6	0	0	7
	14.3	85.7	0.0	0.0	100
	0.0	0.0	0.0	0.0	0.0
Unemployed	1,507	697	145	5	2,354
	64.0	29.6	6.2	0.2	100
	4.3	2.2	3.7	4.8	3.3
Home Duties	1,417	1,031	211	7	2,666
	53.2	38.7	7.9	0.3	100
	4.0	3.2	5.3	6.7	3.7
Not Classifiable	9,340	6,429	1,000	26	16,795
	55.6	38.3	6.0	0.2	100
Not Chated	26.4	20.0	25.3	24.8	23.5
Not Stated	57	81	12	0	150
	38.0	54.0	8.0	0.0	100 0.2
Total	0.2 <b>35,362</b>	0.3 <b>32,152</b>	0.3 <b>3,951</b>	0.0 <b>105</b>	71,570
Total	35,362 49.4	44.9	5.5	0.1	100
	100	100	100	100	100
	100	100	100	100	100

Note: See Appendix C for a description of the classification system for occupation.

FIGURE 4.8 Percentage Distribution of Infants Feeding by Mother's Occupation, Total Live Births, 2007



See Table 4.13 for data. 'Not Stated' values are not included in the calculation of percentages. Notes: Categories where percentages would be based on <100 births are not included in this graph.

# **SECTION FIVE**

# **PERINATAL OUTCOMES**



#### INTRODUCTION

The NPRS data set collects morbidity and mortality data on mothers giving birth and their babies. The Tenth Revision of the International Classification of Diseases (ICD-10)<sup>20</sup> came into effect for coding Irish births from 1 January 2005 onwards. The NPRS coders are trained in ICD-10 coding practices, with particular focus on chapters Pregnancy, childbirth and the puerperium, Certain conditions originating in the perinatal period, and Congenital malformation, deformations and chromosomal abnormalities (Chapters XV - XVII). See Appendix D for a detailed description on cause of death classification and coding rules applied by coders in the NPRS Unit at the ESRI.

This section presents a detailed review on aspects of perinatal outcomes for the infant. Singleton and multiple births are reported in separate tables. Cause of death is examined in greater detail according to the guidelines set out by the WHO in ICD-10. The following tables outline data for live births, antepartum and intrapartum stillbirths, and early neonatal deaths, together with the relevant rates.

Numbers of deaths, mortality rates, and cause of death are presented by birthweight, highlighting percentages of deaths by type of death and birthweight as well as details of rates by cause of death within each birthweight category. Numbers of deaths and mortality rates by gestational age at delivery are also presented. Cause of death is then examined in greater detail, outlining numbers and percentages by type of death. Age at death for early neonatal deaths and post-mortem examination details are also presented in this section.

<sup>&</sup>lt;sup>20</sup> World Health Organization, 2005, *International Statistical Classification of Diseases and Related Health Problems,* 10<sup>th</sup> Revision, Vols. 1-3. Geneva: World Health Organization.

#### **MORTALITY BY BIRTHWEIGHT CATEGORY**

Tables 5.1a and 5.1b summarise live births, antepartum and intrapartum stillbirths, early neonatal deaths, and their relevant mortality rates by birthweight groups. Just over two-thirds of singleton perinatal deaths and more than 75 per cent of multiple perinatal deaths weighed less than 2,500 grams (see Figures 5.1 and 5.2). Low birthweight babies (weighing less than 2,500 grams) are particularly vulnerable and had higher perinatal mortality rates (see Figure 5.3). The perinatal mortality rate for low birthweight babies was 115.8 per 1,000 for singleton births and 19.2 per 1,000 for multiple births. This compares to a perinatal mortality rate of 2.4 per 1,000 singleton and 6.2 per 1,000 multiple births weighing 2,500 grams or more.

**TABLE 5.1a: SINGLETON BIRTHS**Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal Deaths, and Mortality Rates by Birthweight, 2007

Birthweight (grams)	Live Births	Per Cent (%) Live Births	Ante- partum Stillbirths	Intra- partum Stillbirths	Not Stated	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
500-749	86	0.1	58	6	3	437.9	35	407.0	666.7	645.8
750-999	91	0.1	24	1	1	222.2	12	131.9	324.8	261.7
1000-1249	131	0.2	30	0	0	186.3	5	<i>38.2</i>	217.4	160.0
1250-1499	161	0.2	23	2	1	139.0	9	55.9	187.2	105.9
1500-1999	516	0.7	33	4	4	73.6	16	31.0	102.3	58.4
2000-2499	1,645	2.4	44	2	1	27.8	18	10.9	38.4	19.9
2500-2999	7,462	10.8	42	2	1	6.0	10	1.3	7.3	5.5
3000-3499	23,127	33.4	41	4	5	2.2	7	0.3	2.5	2.1
3500-3999	24,462	35.3	21	4	3	1.1	4	0.2	1.3	1.0
4000-4499	9,658	13.9	7	1	2	1.0	2	0.2	1.2	0.9
4500 and Over	1,969	2.8	2	0	0	1.0	0	0.0	1.0	1.0
Not Stated	10	0.0	2	0	0	166.7	0	0.0	166.7	90.9
Total	69,318	100	327	26	21	5.4	118	1.7	7.1	5.1


<sup>&</sup>lt;sup>21</sup> Throughout this section, 'Not Stated' values are not included in the calculation of percentages for discussion.

**TABLE 5.1b: MULTIPLE BIRTHS** 

Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal Deaths, and Mortality Rates by Birthweight, 2007


Birthweight (grams)	Live Births	Per Cent (%) Live Births	Ante- partum Stillbirths	Intra- partum Stillbirths	Not Stated	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
500-749	19	0.8	0	0	1	50.0	7	368.4	400.0	400.0
750-999	41	1.8	4	0	1	108.7	5	122.0	217.4	200.0
1000-1249	48	2.1	0	0	0	0.0	1	20.8	20.8	0.0
1250-1499	62	2.8	0	0	0	0.0	0	0.0	0.0	0.0
1500-1999	301	13.4	1	0	1	6.6	0	0.0	6.6	6.6
2000-2499	665	29.5	1	0	0	1.5	0	0.0	1.5	1.5
2500-2999	766	34.0	3	1	0	5.2	0	0.0	5.2	5.2
3000-3499	304	13.5	2	0	0	6.5	0	0.0	6.5	6.5
3500-3999	44	2.0	0	0	0	0.0	1	22.7	22.7	0.0
4000-4499	2	0.1	0	0	0	0.0	0	0.0	0.0	0.0
Not Stated	0	0.0	4	0	0	1,000.0	0	0.0	1,000.0	1,000.0
Total	2,252	100	15	1	3	8.4	14	6.2	14.5	12.8

**FIGURE 5.1**Deaths by Birthweight Groups, Singleton Births, 2007




Note: See Table 5.1a for data. 'Not Stated' values are not included in the calculation of percentages.

**FIGURE 5.2**Deaths by Birthweight Groups, Multiple Births, 2007



Note: See Table 5.1b for data. 'Not Stated' values are not included in the calculation of percentages.

**FIGURE 5.3**Perinatal Mortality Rates by Birthweight Groups, Total Births, 2007



Note: See Appendix H, Table H18 for data. Rates based on 20 or less observations are not included in this Figure.

# **CAUSE OF DEATH BY BIRTHWEIGHT**

Tables 5.2a and 5.2b show cause of death by each birthweight group for singleton and multiple births respectively. The relevant rate within each birthweight group is also provided in these tables. Over two-thirds of total singleton and multiple perinatal deaths weighed <2,500 grams. The highest percentage (30.2 per cent) of these deaths were due to congenital malformations. This was followed by immaturity (20.6 per cent) and maternal factors (16.7 per cent), and 15.3 per cent recorded cause of death as unexplained (see Figure 5.4). Almost one-third of total singleton and multiple perinatal deaths weighed at least 2,500 grams. Of these, 23 per cent were due to maternal factors, 20 per cent were due to congenital malformation, and 29.7 per cent recorded cause of death as unexplained (see Figure 5.5).

**TABLE 5.2a: SINGLETON BIRTHS** 

Perinatal Mortality: Numbers and Rates, by Birthweight and Cause of Death, 2007

Cause of Death	ICD-10							Birt	hweight (gra	ms)					
	Code		500-749	750-999	1000- 1249	1250- 1499	1500- 1999	2000- 2499	2500- 2999	3000- 3499	3500- 3999	4000- 4499	4500 and Over	Not Stated	Total
Foetus and newborn	P00-P04	Number	18	6	6	6	9	11	14	11	9	2	1	0	93
affected by maternal factors and by complications of pregnancy, labour and delivery		Rate	117.6	51.3	37.3	32.1	16.2	6.5	1.9	0.5	0.4	0.2	0.5	0.0	1.3
Disorders related to	P05-P08	Number	46	9	2	2	2	0	1	0	0	0	0	1	63
length of gestation and foetal growth		Rate	300.7	76.9	12.4	10.7	3.6	0.0	0.1	0.0	0.0	0.0	0.0	83.3	0.9
Birth trauma	P10-P15	Number	1	0	0	0	0	0	0	0	0	0	0	0	1
		Rate	6.5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Respiratory and	P20-P29	Number	12	4	4	2	6	6	7	9	5	1	0	0	56
cardiovascular disorders specific to the perinatal period		Rate	78.4	34.2	24.8	10.7	10.8	3.5	0.9	0.4	0.2	0.1	0.0	0.0	0.8
Infections specific to the	P35-P39	Number	2	0	2	2	1	0	1	1	0	2	0	0	11
perinatal period		Rate	13.1	0.0	12.4	10.7	1.8	0.0	0.1	0.0	0.0	0.2	0.0	0.0	0.2
Haemorrhagic and	P50-P61	Number	1	0	2	0	0	0	3	3	0	0	0	0	9
haematological disorders of foetus and newborn		Rate	6.5	0.0	12.4	0.0	0.0	0.0	0.4	0.1	0.0	0.0	0.0	0.0	0.1
Transitory endocrine and	P70-P74	Number	0	0	0	0	0	0	0	0	0	1	0	0	1
metabolic disorders specific to foetus and newborn		Rate	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.0
Conditions involving the	P80-P83	Number	2	3	0	2	2	2	1	3	0	0	0	0	15
integument and temperature regulation of foetus and newborn		Rate	13.1	25.6	0.0	10.7	3.6	1.2	0.1	0.1	0.0	0.0	0.0	0.0	0.2
Other disorders	P90.0-	Number	0	0	0	0	0	0	1	1	0	2	0	0	4
originating in the perinatal period	P94.9, P96.0- P96.8	Rate	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.0	0.2	0.0	0.0	0.1

Perinatal Mortality: Numbers and Rates, by Birthweight and Cause of Death, 2007


Singleton Births

Cause of Death	ICD-10							Birt	hweight (gra	ms)					
	Code		500-749	750-999	1000- 1249	1250- 1499	1500- 1999	2000- 2499	2500- 2999	3000- 3499	3500- 3999	4000- 4499	4500 and Over	Not Stated	Total
Foetal death of	P95, P96.9	Number	10	5	8	3	11	14	13	21	11	1	1	0	98
unspecified cause, Condition originating in the perinatal period, unspecified		Rate	65.4	42.7	49.7	16.0	19.7	8.3	1.7	0.9	0.4	0.1	0.5	0.0	1.4
Congenital malformations	Q00-Q07	Number	5	3	1	2	5	10	2	1	1	2	0	0	32
of the nervous system		Rate	32.7	25.6	6.2	10.7	9.0	5.9	0.3	0.0	0.0	0.2	0.0	0.0	0.5
Congenital malformations	Q20-Q28	Number	2	0	1	1	2	2	2	1	2	1	0	0	14
of the circulatory system		Rate	13.1	0.0	6.2	5.3	3.6	1.2	0.3	0.0	0.1	0.1	0.0	0.0	0.2
Congenital malformations	Q30-Q34	Number	0	0	0	0	0	1	0	1	1	0	0	0	3
of the respiratory system		Rate	0.0	0.0	0.0	0.0	0.0	0.6	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Other congenital	Q38-Q45	Number	0	0	0	1	0	0	1	1	0	0	0	0	3
malformations of the digestive system		Rate	0.0	0.0	0.0	5.3	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Congenital malformations	Q60-Q64	Number	0	1	0	3	3	3	2	1	0	0	0	0	13
of the urinary system		Rate	0.0	8.5	0.0	16.0	5.4	1.8	0.3	0.0	0.0	0.0	0.0	0.0	0.2
Congenital malformations	Q65-Q79	Number	0	1	0	2	2	4	1	1	1	0	0	0	12
and deformations of the musculoskeletal system		Rate	0.0	8.5	0.0	10.7	3.6	2.4	0.1	0.0	0.0	0.0	0.0	0.0	0.2
Other congenital	Q80-Q89	Number	1	0	1	2	2	3	0	0	1	0	0	0	10
malformations		Rate	6.5	0.0	6.2	10.7	3.6	1.8	0.0	0.0	0.0	0.0	0.0	0.0	0.1
Chromosomal	Q90-Q99	Number	1	5	8	6	12	9	6	2	1	0	0	1	51
abnormalities, not elsewhere classified		Rate	6.5	42.7	49.7	32.1	21.5	5.3	0.8	0.1	0.0	0.0	0.0	83.3	0.7
All other causes	A00.0-	Number	1	1	0	1	0	0	0	0	0	0	0	0	3
	O99.9, R00.0- Z99.9	Rate	6.5	8.5	0.0	5.3	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
All Causes of Perinatal	-	Number	102	38	35	35	57	65	55	57	32	12	2	2	492
Mortality		Rate	666.7	324.8	217.4	187.2	102.3	38.4	7.3	2.5	1.3	1.2	1.0	166.7	7.1

**TABLE 5.2b: MULTIPLE BIRTHS**Perinatal Mortality: Numbers and Rates, by Birthweight and Cause of Death, 2007


Cause of Death	ICD-10 Code						Birthweig	ht (grams)				
			500-749	750-999	1000-1249	1500-1999	2000-2499	2500-2999	3000-3499	3500-3999	Not Stated	Total
Foetus and newborn affected by maternal factors and by	P00-P04	Number Rate	0 0.0	1 21.7	0 0.0	1 3.3	1 1.5	1 1.3	0 0.0	0 0.0	0 0.0	4 1.8
complications of pregnancy, labour and delivery												
Disorders related to	P05-P08	Number	7	5	0	0	0	0	0	0	3	15
length of gestation and foetal growth		Rate	350.0	108.7	0.0	0.0	0.0	0.0	0.0	0.0	750.0	6.6
Respiratory and	P20-P29	Number	0	0	0	0	0	2	1	0	0	3
cardiovascular disorders specific to the perinatal period		Rate	0.0	0.0	0.0	0.0	0.0	2.6	3.3	0.0	0.0	1.3
Haemorrhagic and	P50-P61	Number	1	1	0	0	0	0	0	0	0	2
haematological disorders of foetus and newborn		Rate	50.0	21.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.9
Foetal death of	P95, P96.9	Number	0	2	0	1	0	1	1	0	0	5
unspecified cause, Condition originating in the perinatal period, unspecified		Rate	0.0	43.5	0.0	3.3	0.0	1.3	3.3	0.0	0.0	2.2
Other congenital	Q80-Q89	Number	0	1	0	0	0	0	0	1	0	2
malformations		Rate	0.0	21.7	0.0	0.0	0.0	0.0	0.0	22.7	0.0	0.9
Chromosomal	Q90-Q99	Number	0	0	1	0	0	0	0	0	1	2
abnormalities, not elsewhere classified		Rate	0.0	0.0	20.8	0.0	0.0	0.0	0.0	0.0	250.0	0.9
Total	-	Number	8	10	1	2	1	4	2	1	4	33
		Rate	400.0	217.4	20.8	6.6	1.5	5.2	6.5	22.7	1,000.0	14.5

FIGURE 5.4 Total Deaths < 2,500 grams by Cause of Death, 2007



Note: See Tables 5.2a and 5.2b for data. 'Not Stated' values are not included in the calculation of percentages.

FIGURE 5.5 Total Deaths ≥ 2,500 grams by Cause of Death, 2007



Note: See note to Figure 5.4.

#### **MORTALITY BY GESTATIONAL AGE**

Tables 5.3a and 5.3b illustrate the significant impact of prematurity (less than 37 completed weeks gestation) on stillbirth and early neonatal mortality rates. Similar to low birthweight babies, almost two-thirds of singleton perinatal deaths were born preterm while 75.7 per cent of multiple perinatal deaths were preterm (see Figures 5.6 and 5.7). The perinatal mortality rates for babies born before 37 completed weeks of gestation were 97.6 per 1,000 singleton births and 22.8 per 1,000 multiple births. For babies born at 37 or more completed weeks these rates were 2.6 per 1,000 singleton births and 6.8 per 1,000 multiple births. Figure 5.8 shows the perinatal mortality rates by gestational age at delivery for total births, illustrating the higher rates associated with lower gestational age.

**TABLE 5.3a: SINGLETON BIRTHS**Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal Deaths, and Mortality Rates by Gestational Age, 2007


Gestational Age at Delivery (weeks)	Live Births	Per Cent (%) Live Births	Ante- partum Stillbirths	Intra- partum Stillbirths	Not Stated	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Less than 22	1	0.0	4	0	1	833.3	1	1,000.0	1,000.0	1,000.0
22-27	163	0.2	60	7	3	300.4	41	251.5	476.4	452.9
28-31	368	0.5	62	0	2	148.1	16	43.5	185.2	139.4
32-36	2,535	3.7	76	4	4	32.1	40	15.8	47.3	24.2
37-41	63,784	92.0	122	15	11	2.3	20	0.3	2.6	2.0
42 and Over	2,457	3.5	3	0	0	1.2	0	0.0	1.2	0.8
Not Stated	10	0.0	0	0	0	0.0	0	0.0	0.0	0.0
Total	69,318	100	327	26	21	5.4	118	1.7	7.1	5.1

**TABLE 5.3b: MULTIPLE BIRTHS** 

Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal Deaths, and Mortality Rates by Gestational Age, 2007


Gestational Age at Delivery (weeks)	Live Births	Per Cent (%) Live Births	Ante- partum Stillbirths	Intra- partum Stillbirths	Not Stated	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Less than 22	0	0.0	2	0	0	1000.0	0	0.0	1,000.0	1,000.0
22-27	49	2.2	1	0	1	39.2	12	244.9	274.5	260.0
28-31	147	6.5	1	0	0	6.8	0	0.0	6.8	6.8
32-36	887	39.4	5	0	2	7.8	1	1.1	8.9	7.8
37-41	1,167	51.8	6	1	0	6.0	1	0.9	6.8	5.1
42 and Over	2	0.1	0	0	0	0.0	0	0.0	0.0	0.0
Total	2,252	100	15	1	3	8.4	14	6.2	14.5	12.8

FIGURE 5.6 Deaths by Gestational Age Groups, Singleton Births, 2007




Note: See Table 5.3a for data.

**FIGURE 5.7**Deaths by Gestational Age Groups, Multiple Births, 2007



Note: See Table 5.3b for data.

**FIGURE 5.8**Perinatal Mortality Rates by Gestational Age at Delivery, Total Births, 2007



Note: See Appendix H, Table H19 for data. Rates based on 20 or less observations are not included in this Figure.

#### **CAUSE OF DEATH**

Of the 71,963 births recorded by the NPRS in 2007, 393 were stillbirths and 132 were early neonatal deaths, giving a total of 525 perinatal deaths and a perinatal mortality rate of 7.3 per 1,000 live births and stillbirths. As noted in Section Two, although the Irish perinatal mortality rate has been generally declining, it is still relatively high in comparison to other EU member states. Meaningful international and European comparisons by cause of death, however, are complicated by differences across countries relating to classifications, data availability, and definition of perinatal death. Some countries report on cause of death for stillbirths only, <sup>22</sup> while others include late neonatal deaths (7 to 27 days) in their perinatal statistics.<sup>23</sup> The European Perinatal Health Report compiled from the EURO-PERISTAT perinatal health indicators compares rates of perinatal mortality across EU countries for 2004 and compares cause of foetal death due to congenital anomalies only. The indicator for causes of all perinatal deaths has been highlighted as an indicator for future development in this project.<sup>24</sup>

Since 2005, mortality data in the NPRS data set has been coded using the WHO ICD-10 classification.<sup>25</sup> Under this system, the main cause of death is defined as the pathological condition of the infant or foetus, which in the opinion of the certifier, made the greatest contribution towards the death.<sup>26</sup> Each year around 20 per cent of all perinatal deaths are classified as 'unspecified/unexplained'. With approximately half of all perinatal deaths in Ireland not undergoing post mortem, it is perhaps unsurprising that such a high percentage of deaths are classified as such. While this poses limitations in the application of the data for clinicians, epidemiologists and service planners, it is, however, not a situation unique to Ireland, as many countries report high percentages of unexplained deaths in the perinatal period in their perinatal data sets, even where different classification systems are used. The UK Confidential Enquiry into Maternal and Child Health (CEMACH), for example, employs the Wigglesworth Classification and for 2007 reported unknown as cause of death for 76.1 per cent of stillbirths and 1.4 per cent of neonatal deaths.<sup>27</sup>

<sup>&</sup>lt;sup>22</sup> See, for example, Canadian Perinatal Health Report, 2008 Edition. Public Health Agency of Canada. Canadian Perinatal Health Report, 2008 Edition. Ottawa, 2008. http://www.publichealth.gc.ca/cphr/

<sup>&</sup>lt;sup>3</sup> See, for example, *Australia's Mothers and Babies, 2006.* Laws, P.J. & Hilder ,L. 2008. Australia's mothers and babies 2006. Perinatal statistics series no. 22. Cat. No. PER 46. Sydney: AIHW National Perinatal Statistics Unit.

EURO-PERISTAT Project, with SCPE, EUROCAT, EURONEOSTAT. European Perinatal Health Report. 2008. www.europeristat.com

World Health Organisation, 2005, International Statistical Classification of Diseases and Related Health Problems, 10<sup>th</sup> Revision, Vols. 1-3. Geneva: World Health Organisation.

<sup>&</sup>lt;sup>26</sup> See Appendix D for more detail

<sup>&</sup>lt;sup>27</sup> Confidential Enquiry into Maternal and Child Health (CEMACH) Perinatal Mortality 2007: United Kingdom. CEMACH: London, 2009.

**TABLE 5.4a: SINGLETON BIRTHS** 

Stillbirths, Early Neonatal Deaths and Mortality Rates by Cause of Death by ICD-10 Chapter and Selected Sub-Categories, 2007

Cause of Death	ICD-10 Code	Stillbirths	Stillbirth	Early	Early	Total	Perinatal
			Rate	Neonatal Deaths	Neonatal Mortality Rate	Perinatal Deaths	Mortality Rate
Foetus and newborn affected by	P00-P04	90	1.3	3	0.0	93	1.3
maternal factors and by complications of pregnancy, labour and delivery				-			
Foetus and newborn affected by maternal conditions that may be unrelated to present pregnancy	P00.0 - P00.9	5	0.1	0	0.0	5	0.1
Foetus and newborn affected by maternal complications of pregnancy	P01.0 - P01.9	6	0.1	0	0.0	6	0.1
Foetus and newborn affected by complications of placenta, cord and membranes	P02.0 - P02.9	69	1.0	3	0.0	72	1.0
Foetus and newborn affected by other complications of labour and delivery	P03.0 - P03.9	9	0.1	0	0.0	9	0.1
Disorders related to length of gestation and foetal growth	P05-P08	30	0.4	33	0.5	63	0.9
Slow foetal growth and foetal malnutrition	P05.0 - P05.9	8	0.1	0	0.0	8	0.1
Disorders related to short gestation and low birthweight, not elsewhere classified	P07.0 - P07.3	22	0.3	33	0.5	55	0.8
Birth trauma	P10-P15	1	0.0	0	0.0	1	0.0
Respiratory and cardiovascular disorders specific to the perinatal period	P20-P29	40	0.6	16	0.2	56	0.8
Intrauterine hypoxia	P20.0 - P20.9	38	0.5	0	0.0	38	0.5
Other respiratory conditions originating in the perinatal Period	P28.0 - P28.9	0	0.0	8	0.1	8	0.1
Infections specific to the perinatal period	P35-P39	9	0.1	2	0.0	11	0.2
Bacterial sepsis of newborn	P36.0 - P36.9	5	0.1	1	0.0	6	0.1
Other infections specific to the perinatal period	P39.0 - P39.9	4	0.1	1	0.0	5	0.1
Haemorrhagic and haematological disorders of foetus and newborn	P50-P61	7	0.1	2	0.0	9	0.1
Foetal blood loss	P50.0 - P50.9	3	0.0	1	0.0	4	0.1
Intracranial nontraumatic haemorrhage of foetus and newborn	P52.0 - P52.9	3	0.0	1	0.0	4	0.1
Transitory endocrine and metabolic disorders specific to foetus and newborn	P70-P74	1	0.0	0	0.0	1	0.0
Conditions involving the integument and temperature regulation of foetus and newborn	P80-P83	11	0.2	4	0.1	15	0.2
Other disorders originating in the perinatal period	P90-P96	97	1.4	5	0.1	102	1.5
Other disturbances of cerebral status of newborn	P91.0-P91.9	1	0.0	3	0.0	4	0.1
Foetal death of unspecified cause, Condition originating in the perinatal period, unspecified	P95, P96.9	96	1.4	2	0.0	98	1.4

**TABLE 5.4a: SINGLETON BIRTHS (contd.)** 

Stillbirths, Early Neonatal Deaths and Mortality Rates by Cause of Death by ICD-10 Chapter and Selected Sub-Categories, 2007

Cause of Death	ICD-10 Code	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Total Perinatal Deaths	Perinatal Mortality Rate
Congenital malformations of the nervous system	Q00-Q07	17	0.2	15	0.2	32	0.5
Anencephaly and similar malformations	Q00.0 - Q00.2	14	0.2	8	0.1	22	0.3
Congenital hydrocephalus	Q03.0 - Q03.9	1	0.0	3	0.0	4	0.1
Other congenital malformations of the brain	Q04.0 - Q04.9	1	0.0	3	0.0	4	0.1
Congenital malformations of the circulatory system	Q20-Q28	9	0.1	5	0.1	14	0.2
Other congenital malformations of heart	Q24.0 - Q24.9	3	0.0	3	0.0	6	0.1
Congenital malformations of the respiratory system	Q30-Q34	2	0.0	1	0.0	3	0.0
Other congenital malformations of the digestive system	Q38-Q45	1	0.0	2	0.0	3	0.0
Congenital malformations of the urinary system	Q60-Q64	4	0.1	9	0.1	13	0.2
Renal agenesis and other reduction defects of kidney	Q60.0 - Q60.6	2	0.0	6	0.1	8	0.1
Congenital malformations and deformations of the musculoskeletal system	Q65-Q79	5	0.1	7	0.1	12	0.2
Congenital malformations of musculoskeletal system, not elsewhere classified	Q79.0 - Q79.9	3	0.0	2	0.0	5	0.1
Other congenital malformations	Q80-Q89	9	0.1	1	0.0	10	0.1
Chromosomal abnormalities, not elsewhere classified	Q90-Q99	38	0.5	13	0.2	51	0.7
Down's syndrome	Q90.0 - Q90.9	11	0.2	1	0.0	12	0.2
Edward's syndrome and Patau's syndrome	Q91.0 - Q91.7	16	0.2	12	0.2	28	0.4
Other trisomies and partial trisomies of the autosomes, not elsewhere classified	Q92.0 - Q92.9	6	0.1	0	0.0	6	0.1
All other causes	A00.0-O99.9, R00.0-Z99.9	3	0.0	0	0.0	3	0.0
Total	-	374	5.4	118	1.7	492	7.1

**TABLE 5.4b: MULTIPLE BIRTHS** 


Stillbirths, Early Neonatal Deaths and Mortality Rates by Cause of Death by ICD-10 Chapter, 2007

Cause of Death	ICD-10 Code	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Total Perinatal Deaths	Perinatal Mortality Rate
Foetus and newborn affected by maternal factors and by complications of pregnancy, labour and delivery	P00-P04	4	1.8	0	0.0	4	1.8
Disorders related to length of gestation and foetal growth	P05-P08	6	2.6	9	4.0	15	6.6
Respiratory and cardiovascular disorders specific to the perinatal period	P20-P29	3	1.3	0	0.0	3	1.3
Haemorrhagic and haematological disorders of foetus and newborn	P50-P61	0	0.0	2	0.9	2	0.9
Other disorders originating in the perinatal period	P90-P96	5	2.2	0	0.0	5	2.2
Other congenital malformations	Q80-Q89	0	0.0	2	0.9	2	0.9
Chromosomal abnormalities, not elsewhere classified	Q90-Q99	1	0.4	1	0.4	2	0.9
Total	_	19	8.4	14	6.2	33	14.5

#### **STILLBIRTHS**


Apart from unspecified/unexplained deaths, the leading cause of death for stillbirths was maternal factors (P00-P04) for singleton births (24.1 per cent) and immaturity (P05-P08) for multiple births (31.6 per cent), (see Figures 5.9 and 5.10). The cause of death was unexplained for around one-quarter of all stillbirths in 2007.

FIGURE 5.9 Stillbirths by Cause of Death, Singleton Births, 2007



Note: See Table 5.4a for data.

**FIGURE 5.10** Stillbirths by Cause of Death, Multiple Births, 2007




Note: See Table 5.4b for data.

#### **EARLY NEONATAL DEATHS**


Almost 45 per cent of singleton early neonatal deaths were due to congenital malformations (Q00-Q99) and a further 28 per cent were due to immaturity (see Figure 5.11). Almost two-thirds of multiple early neonatal deaths were due to immaturity (64.3 per cent) and a further 21.4 per cent were due to congenital malformations (see Figure 5.12).

**FIGURE 5.11**Early Neonatal Deaths, by Cause of Death, Singleton Births, 2007



Note: See Table 5.4a for data. ENNDs denotes Early Neonatal Deaths.

**FIGURE 5.12**Early Neonatal Deaths, by Cause of Death, Multiple Births, 2007




Note: See Table 5.4b for data. ENNDs denotes Early Neonatal Deaths.

# **TOTAL PERINATAL DEATHS**


The leading cause of death for total perinatal deaths was congenital malformations (27 per cent) followed by maternal factors (18.5 per cent). For singleton births, congenital malformations were the greatest contributor to cause of death, accounting for 28 per cent of total singleton perinatal deaths (see Figure 5.13). For multiple births, the largest contributor was immaturity (45.5 per cent) followed by maternal factors and congenital malformations, 12.1 per cent in each (see Figure 5.14). Overall, 19.9 per cent and 15.2 per cent of singleton and multiple deaths were unexplained.

**FIGURE 5.13**Perinatal Deaths by Cause of Death, Singleton Births, 2007



Note: See Table 5.4a for data.

FIGURE 5.14
Perinatal Deaths by Cause of Death, Multiple Births, 2007



Note: See Table 5.4b for data.

### **AGE AT DEATH**

Age at death is recorded for early neonatal deaths and is shown in Tables 5.5a and 5.5b. Just over 35 per cent of total early neonatal deaths happened within the first hour of life. A total of 73.6 per cent of early neonatal deaths occurred before completing the first day of life and a further 26.4 per cent occurred before completing the first week of life.

**TABLE 5.5a: SINGLETON BIRTHS**Age at Death, Early Neonatal Deaths, 2007

Time	Frequency	Per Cent (%)
Completed Hours		
<1	42	35.6
1	20	16.9
2	9	7.6
3	3	2.5
4	2	1.7
5	1	0.8
6	2	1.7
7	1	0.8
11	1	0.8
12	3	2.5
13	1	0.8
15	1	0.8
17	1	0.8
19	1	0.8
20	1	0.8
22	1	0.8
Total under 1 day	90	76.3
<b>Completed Days</b>		
1	12	10.2
2	1	0.8
3	3	2.5
4	6	5.1
5	1	0.8
6	2	1.7
Total under 1 Week	25	21.2
Unknown	3	2.5
Total	118	100

**TABLE 5.5b: MULTIPLE BIRTHS** 

Age at Death, Early Neonatal Deaths, 2007

Time	Frequency	Per Cent (%)
<b>Completed Hours</b>		
<1	4	28.6
1	1	7.1
Total under 1 day	5	35.7
<b>Completed Days</b>		
1	2	14.3
2	3	21.4
3	2	14.3
5	2	14.3
Total under 1 Week	9	64.3
Total	14	100

### **POST-MORTEM EXAMINATION UPTAKE**

Data on autopsy uptake are shown in Tables 5.6a and 5.6b. Post-mortems were less likely in the case of early neonatal deaths compared to stillbirths and this trend is consistent over recent years. The percentage of early neonatal deaths undergoing post-mortem examinations was 30.8 per cent compared to 48.3 per cent for stillbirths. An autopsy was undertaken in 44 per cent of total perinatal deaths in 2007.

**TABLE 5.6a: SINGLETON BIRTHS** Stillbirths and Early Neonatal Deaths undergoing Post-Mortem Examinations, 2007

Post Mortem		Type of Death	
Frequency	Early Neonatal	Stillbirths	Total
Row Per Cent	Deaths		
Col. Per Cent			
Post-Mortem	35	180	215
	16.3	83.7	100
	29.7	48.1	43.7
No Post-Mortem	81	192	273
	29.7	70.3	100
	68.6	51.3	55.5
Not Stated	2	2	4
	50.0	50.0	100
	1.7	0.5	0.8
Total	118	374	492
	24.0	76.0	100
	100	100	100

**TABLE 5.6b: MULTIPLE BIRTHS** 

Stillbirths and Early Neonatal Deaths undergoing Post-Mortem Examinations, 2007

Post Mortem		Type of Death						
Frequency	Early Neonatal	Stillbirths	Total					
Row Per Cent	Deaths							
Col. Per Cent								
Post-Mortem	5	9	14					
	35.7	64.3	100					
	35.7	47.4	42.4					
No Post-Mortem	9	10	19					
	47.4	52.6	100					
	64.3	52.6	57.6					
Total	14	19	33					
	42.4	57.6	100					
	100	100	100					

# **APPENDICES**



# **APPENDIX A**

# **DEFINITIONS**

This section collates the definitions of the terms used in the analyses for this report.

# **OUTCOMES AND RATES<sup>28</sup>**

Perinatal Period: The perinatal period commences at 22 completed weeks (154 days) of gestation (the time when birth weight is normally 500 g), and ends seven completed days after birth.

Birth Rate: The Birth Rate or Crude Birth Rate is the ratio of total live births to total population over a specified period of time. The birth rate is often expressed as the number of live births per 1,000 members of the population in a particular time period (for example year).

Live Birth: A live birth is defined as the complete expulsion or extraction from its mother of a product of conception, irrespective of the duration of pregnancy, which, after such separation, breathes or shows any other evidence of life, such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles, whether or not the umbilical cord has been cut or the placenta is attached. In accordance with WHO guidelines, live births weighing less than 500 grams are not included in the national statistics presented in this report.

Stillbirth and Stillbirth Rate: Foetal death is defined as death prior to the complete expulsion or extraction from its mother of a product of conception, irrespective of the duration of the pregnancy. A foetal death is indicated by the fact that, after such separation, the foetus does not breathe or show any other evidence of life, such as beating of the heart, pulsation of the umbilical cord, or definite movement of voluntary muscles. Following WHO guidelines, a stillbirth in this report, refers to the death of a foetus weighing at least 500 grams. The stillbirth rate is given as follows:

> Number of Stillbirths x 1,000 Total Number of Live Births and Stillbirths

<sup>&</sup>lt;sup>28</sup> In this report, where rates are based on very small numbers, they should be interpreted with caution. Rates calculated from 20 cases or less are distinguished by italic type as a warning to the user that their reliability as a measure may be affected by the small number of observations.

Number of Early Neonatal Deaths x 1,000

Total Number of Live Births

Perinatal Death and Perinatal Mortality Rate (PMR): Perinatal deaths include stillbirths and early neonatal deaths. The perinatal mortality rate (PMR) is calculated as follows:

Number of Stillbirths and Early Neonatal Deaths x 1,000

Total Number of Live Births and Stillbirths

In this report a second perinatal mortality rate has been calculated excluding all stillbirths and early neonatal deaths due to congenital anomalies. This adjusted rate (Adjusted PMR) to some extent removes those perinatal events where death was unavoidable, but no attempt has been made to exclude only invariably fatal congenital anomalies.

Congenital Anomalies are physiological or structural abnormalities that develop at or before birth and are present at the time of birth. Congenital anomalies are those diseases/conditions categorised in *ICD-10*, Chapter XVII, *Congenital Malformations*, *Deformities and Chromosomal Abnormalities* (Q00.0-Q99.9).

Twinning Rate: This is given by:

Number of Twin Maternities x 1,000

Total Number of Maternities

Twin maternities, which resulted in stillbirths, are included in the calculation of the twinning rate.

# **O**THER ITEMS

The definitions given below relate to the principal classification variables contained in the main section of the report and refer only to those terms where explanation is considered necessary. The definitions are grouped according to the section in which they first appear.

#### **GENERAL CHARACTERISTICS**

Born Before Arrival (BBA): Born before arrival indicates that an infant was delivered before arrival at hospital.

Domiciliary Birth: Domiciliary births are planned home births. The vast majority of these are attended by an independent midwife and are not associated with a hospital. In 2007, there were 46 births under hospital administered home birth schemes. For the purposes of this report these births are treated as hospital births due to their small numbers.

Marital Status: Five categories are recognised as specified below:

- 1. Married
- 2. Never Married<sup>29</sup>
- 3. Widowed
- 4. Separated, covering the following categories:
  - Deserted
  - Legally separated
  - Church decree of nullity
  - Otherwise separated
- 5. Divorced.

Mother's and Father's Occupation: Occupations are coded and grouped, with minor modifications, according to the system of socio-economic groupings as used by the Central Statistics Office in the 1991 Census of Population, Volume 6. Details are given in Appendix C.

Mother's and Father's Nationality: This indicates place of birth for mother and father. This definition is for the sole purpose of giving the necessary directions for the completion of the Birth Notification Form and is not a legal definition.

Parity: Maternal parity indicates a mother's total number of previous live births and stillbirths.

Time Since Last Birth: This variable gives the time, grouped into yearly intervals, since the mother's last live birth or stillbirth.

Birthweight: Birthweight is given in grams and hospitals are asked to record the weight within the first hour after birth. In accordance with WHO guidelines, infants weighing under 500 grams are not included in the national statistics presented in this report.

Period of Gestation: Gestational age at delivery is recorded in completed weeks, and is measured or estimated according to the best method available in each case.

<sup>&</sup>lt;sup>29</sup> Data recording marital status as 'Never Married' are classified as 'Single' marital status in discussion in this report.

Antenatal Length of Stay: This is computed as the difference, in days, between a mother's date of admission and her date of delivery. Domiciliary births are not included in tables relating to this variable.

Postnatal Length of Stay: This is computed as the difference, in days, between the date of delivery and the date of the mother's discharge from hospital. A return of 'not stated' for this variable may indicate that the completed form was returned before the mother had been discharged. Domiciliary births are not included in tables relating to this variable.

Hospital Admission Booked or Unbooked: This indicates whether a mother's admission into hospital for delivery was booked or unbooked prior to delivery. Domiciliary births are not included in the tables relating to this variable.

Immunity to Rubella: An affirmative answer should be recorded only where there is documented evidence of a rubella antibody test. A history of rubella or previous immunisation is not sufficient.

Method of Delivery: Six categories are used:

- 1. Spontaneous cephalic delivery
- 2. Breech delivery, spontaneous, with or without forceps to after-coming head, breech extraction
- 3. Forceps delivery
- 4. Vacuum extraction<sup>30</sup>
- 5. Caesarean section
- 6. Other Specified and Combined (for example, combination of spontaneous delivery followed by a vacuum extraction or caesarean section).

Size of Maternity Unit: Size in this context refers to the number of live births and stillbirths occurring during the year.

Transfer of Infants: This identifies infants who were transferred to another hospital for medical reasons.

Length of Infant's Stay: This is computed as the difference, in days, between the infant's date of birth and the date of the infant's discharge from hospital. A return of 'not stated' for this variable may indicate that the completed form was returned before the infant had been discharged. Where an infant was transferred to another hospital, the date of the transfer is

<sup>&</sup>lt;sup>30</sup> Also known as Ventouse Deliveries (*Source*: World Health Organization, 2005, *International Statistical Classification of Diseases and Related Health Problems*, 10<sup>th</sup> Revision, Vol. 1. Geneva: World Health Organization.)

given as the date of the infant's discharge. Domiciliary births and perinatal deaths are not included in the tables relating to this variable.

Type of Feeding: This indicates whether artificial feeding, breastfeeding, or a combination of both was being used at the time of discharge. The Breastfeeding Rate is based on live births only (excluding ENNDs), and is calculated as follows:

> Number of Mothers Breastfeeding x 100 Total Number of Live Births (excluding ENNDs)

#### **PERINATAL OUTCOMES**

Antepartum and Intrapartum Stillbirths: These refer to stillbirths occurring prior to and during labour respectively.

Cause of Death: This categorisation indicates the principal underlying pathological cause, which in the opinion of the certifier made the greatest contribution to the death of the infant or foetus. In assigning cause of death in the NPRS, the guidelines recommended by the WHO have been followed. A summary of these guidelines is provided in Appendix D. The Tenth Revision of the ICD (ICD-10) came into effect for coding Irish births from 1 January 2005 onwards.

Age at Death: The age of early neonatal deaths is given in completed hours if the infant survived less than one day and in completed days otherwise.

Post-mortem: This indicates whether or not a post-mortem examination was performed. In some cases a return of 'not stated' may be due to the notification form being returned prior to post-mortem.

Note: All perinatal statistics and definitions are based on WHO ICD-10 Classifications.

# **APPENDIX B**

# **COPY OF PART 3 OF THE BIRTH NOTIFICATION FORM USED BY NPRS**

The following format was used for births in 2007.

	-			e, 11mmmen 0 quant, 00	r John Rogerson's Quay, Dublin 2
TYPE OF BERTH 1 LL Live = 1, Sell = 2)	PLACE OF BIRTH (III	opital = 1, BBA = 2, Domicitary = 3)	2 🔲	NAME AND	
1. Sti = 2)	HOSPITAL NO. 3	CASE	1	ADDRESS OF	
	NO. 3	NO. 6	YYY	HOSPITAL	
DATE OF BIRTH (DD/MM/YYYY)	14		MOTHER'S HEALTH		
	TIME OF BIRTH		ANTENATAL CARE THIS PR (Bospital / Obstatrician = 1, G)		ne = 4. Midwife Onlys5) 117
IF MULTIPLE BIRTH	ORDER OF BUILTH	No. 22 of 23			
			DATE OF FIRST VISIT TO DO DURING PREGNANCY (DD)		118
			DATE OF FIRST VISIT TO H		
SEX (Male = 1, Female = 2, Indetermin	ate = 3)	24	DURING PREGNANCY (DD)		126
BUCHWEIGHT		25 GRAMMES	WAS MOTHER IMMUNE TO	RUBELLA	
PERSOD OF GESTATION		29 WEBKS	(Yes = 1, No = 2, Not Known :		134
FATHER'S DETAILS			METHOD OF DELIVERY (Sp	ontaneous = 1. Breach + Force	ns = 2.
			Forceps = 3, Vac. Extraction =	4, Caesseun Sec. = 5, Other =	6) 135
			MAIN MATERNAL DESEASE	OR CONDITION A PRICTIN	IC FORTIS OR IN EAST
			MALE MALESTAL DISEASE	OR COMBINEDAMENT	D POR CO OR ENTANT
					136
			OTHER MATERNAL DESEAS	ES OR CONDITIONS AFFEC	TING POETUS OR INFANT
COUNTY		.1 1			
		31		-	1
					141
NATIONALITY		38 42 1	INFANT'S HEALTH		1
DATE OF BIRTH (DDMMYYYY)			TYPE OF PEEDING (Artificial	l = 1. Breast = 2. Combined = :	3) 146
DATE OF BURIET (DUMMYYYY)	44 L		WAS BCG ADMINISTERED (	Yea = 1, No = 2)	147
MOTHER'S DETAILS			MAIN DESEASE OR CONCID	NITAL MALFORMATION AF	FECTING INFANT
					148
			OTHER DESEASES OR COING	ZENITAL MALFORMATION	APPECTING IN FANT
					153
			HOSPITAL		148
COUNTY		52	WAS ADMESSION BOOK ED	(Yea = 1, No = 2)	156
			DATE OF MOTHER'S ADMIS	SION (DDMMYYYY)	159
NATIONALITY			DATE OF MOTHER'S DISCH	ARGE (DOMM YYYY)	167
OCCUPATION		63			
DATE OF BIRTH (DOMMYYYY)	65		DATE OF INBANT'S DISCHA	IRGE (DDMMYYYY)	175
			WAS INSANT TRANSFERRE	D TO OTHER HOSPITAL	
MARITAL STATUS (Market = 1, News M	aried = 1, Wilcord = 1, Marie dea	Separated = 4,Discovering = 5) 73	FOR MEDICAL REASONS (1	(a = 1, No = 2)	183
DATE OF PRESENT MARRIAGE (DE	MMYYYY) 74		IF 'YES', NAME OF HOSPIT	AL	
DATE OF LAST BURTH (by or still) (			and the second		
	EBERTHS	90 📖			
					184
	LDREN STILL LIVING	92	GENERAL PRACTITIO	NER ATTENDED BY M	OTHER
	LBIRTHS	» LL			
PERINATAL DEATH	NTANEOUS ABORTIONS				
TYPE OF DEATH (Early Necessal = 1. WAS AUTOPSY PERFORMED (Va. =		98			
WAS AUTOPSY PERFORMED (SEE = AGE AT DEATH	1.No=2) 100 DAYS	101 LL HOURS			
		1			
PLACE OF DEATH	OCCUR BEFORE LABOUR	(1) 103			
	BOUR (2) NOT KNOWN (5)	106			
CAUSE OF DEATH MAIN DISEASE OR CONDITIONIN	FOETUS OR INFANT.				
		107			
OTHER DISEASES OR CONDITIONS	IN POETUS OR IN FAINT _				
		112			
Signature		Dute			BNF01/2003

## **APPENDIX C**

#### CLASSIFACTION OF OCCUPATION

Occupations have been coded and classified in a set of socio-economic groups according to a system devised by the CSO. The final CSO Category of 'Unknown' has been differentiated into five distinct groups for the purposes of the Perinatal Reporting System. These are 'Unemployed', 'Not Classifiable', 'Not Applicable', 'Home Duties', and 'Not Stated'. Occupation is coded as unemployed when occupation is given as unemployed and where no previous occupation is stated. If a previous occupation is stated then occupation is coded to the relevant category. 'Not Classifiable' has been used to categorise indecipherable, unclear, or unlisted occupations where efforts to clarify the information have failed. 'Home Duties' is entered where occupation has been entered as 'full-time mother/father/parent', 'stay at home mother/father/parent', 'housewife', 'home duties', etc. 'Not Stated' applies to those cases where either the mother's or father's occupation has been left blank or has been recorded as 'unknown'.

#### Under Father's occupation, the following applies:

'Not Applicable' is entered when the marital status of the mother has been given as never married, widowed, separated or divorced, and where father's occupation has been left blank. 'Not Stated' applies to those cases where the marital status of the mother is recorded as either married or not stated, and where father's occupation has been left blank or has been recorded as 'unknown'. In either case, if an occupation is given for the father then the appropriate occupational code has been used.

#### The full list of socio-economic groups used in this report is as follows:

- Farmers & Farm Managers
- Other Agricultural Occupations & Fishermen
- **Higher Professional**
- **Lower Professional**
- **Employers & Managers**
- Salaried Employees
- Intermediate Non-Manual Workers
- Other Non-Manual Workers

- **Skilled Manual Workers**
- Semi-Skilled Manual Workers
- **Unskilled Manual Workers**
- Unemployed
- Not Classifiable
- Not Applicable (Father's Occupation Only)
- **Home Duties**
- Not Stated.

## **APPENDIX D**

#### **CAUSE OF DEATH CLASSIFICATION**

The main cause of death has been determined according to the guidelines set out by the WHO in the Tenth Revision of the ICD.<sup>31</sup> Briefly, the principal coding criteria are as follows:

The main cause of death as far as possible should indicate the pathological condition of the infant or foetus, which in the opinion of the certifier made the greatest contribution towards the death.

The following rules are extracted from *ICD-10*, *Second Edition*, *2005*, *4*. *Rules and guidelines for mortality and morbidity coding*. These rules have been applied to cause of death coding in the NPRS data set as follows:

Section (A) – main disease or condition in foetus or infant

Section (B) – other diseases or conditions in foetus or infant

Section (C) – main maternal disease or condition affecting foetus or infant

Section (D) – other maternal disease or condition affecting foetus or infant.

#### Rule P1. Mode of death or prematurity entered in section (a)

If heart or cardiac failure, asphyxia or anoxia (any condition in P20.-, P21.-) or prematurity (any condition in P07.-) is entered in section (a) and other conditions of the infant or foetus are entered in either section (a) or section (b), code the first-mentioned of these other conditions as if it had been entered alone in section (a) and code the condition actually entered in section (a) as if it had been entered in section (b).

#### Rule P2. Two or more conditions entered in sections (a) or (c)

If two or more conditions are entered in section (a) or section (c), code the first-mentioned of these as if it had been entered alone in section (a) or (c) and code the others as if they had been entered in sections (b) or (d).

<sup>&</sup>lt;sup>31</sup> World Health Organization, 2005, *International Statistical Classification of Diseases and Related Health Problems*, 10<sup>th</sup> Revision, Vols. 1-3. Geneva: World Health Organization.

#### Rule P3. No entry in sections (a) or (c).

If there is no entry in section (a) but there are conditions of the infant or foetus entered in section (b), code the first-mentioned of these as if it had been entered in section (a); if there are no entries in either section (a) or section (b), either code P95 (Foetal death of unspecified cause) for stillbirths or code P96.9 (Condition originating in the perinatal period, unspecified) for early neonatal deaths should be used for section (a).

Similarly, if there is no entry in section (c) but there are maternal conditions entered in section (d), code the first-mentioned of these as if it had been entered in section (c); if there are no entries in either section (c) or section (d) use some artificial code, e.g. xxx.x for section (c) to indicate that no maternal condition was reported.

#### Rule P4. Conditions entered in wrong section.

If a maternal condition (i.e. conditions in P00-P04) is entered in section (a) or section (b), or if a condition of the infant or foetus is entered in section (c) or section (d), code the conditions as if they had been entered in the respective correct section.

For the purpose of the cause of death tables included in this report, one further criterion has been applied. Where no infant's condition is given then, where a specific maternal condition is given (any condition in P00-P04), the main cause of death is classified under the maternal condition.

Table D1 summarises these rules.

**TABLE D1**Cause of Death Coding Rules

ICD-10 Rule		Cause of Death as recorded of	on the Birth Notification Form		Action
	Section A	Section B	Section C	Section D	
	Main disease or condition in foetus or infant	Other disease or condition in foetus or infant	Main maternal disease or condition affecting foetus or	Other maternal disease or condition affecting foetus or	
			infant	infant	
Rule 1	Heart or cardiac failure, asphyxia, anoxia, prematurity (P20, P21 or P07)	Other condition	No entry	No entry	Code the stated 'other condition' in section A and move condition entered in section A to section B.
Rule 2	Two or more conditions listed	No entry	Two or more conditions listed	No entry	Code the first mentioned condition in sections A and C respectively and code the second mentioned conditions in sections B and D respectively.
Rule 3	No entry	Condition entered	No entry	Condition entered	Code the first mentioned condition in section B to section A and the first mentioned condition in section D to section C.
	No entry	No entry	No entry	No entry	Code 'Foetal death of unspecified cause' (P95) for stillbirths or 'Conditions originating in the perinatal period, unspecified' (P96.9) for early neonatal deaths in section A.
Rule 4	Maternal condition entered (P00-P04)	Maternal condition entered (P00-P04)	Infant condition entered	Infant condition entered	Code maternal conditions in sections C and D respectively. Code infant conditions in sections A and B respectively.
Additional Rule	No entry	No entry	Maternal condition (P00-P04)	No entry	Code maternal condition to section A.

Table D2 illustrates the main chapter blocks relating to cause of death categories in this report.

**TABLE D2** ICD 10, Cause of Death Codes

ICD-10	CAUSE OF DEATH
CODE	
P00-P04	Foetus and newborn affected by maternal factors and by complications of pregnancy, labour and delivery
P05-P08	Disorders related to length of gestation and foetal growth
P10-P15	Birth trauma
P20-P29	Respiratory and cardiovascular disorders specific to the perinatal period
P35-P39	Infections specific to the perinatal period
P50-P61	Haemorrhagic and haematological disorders of foetus and newborn
P70-P74	Transitory endocrine and metabolic disorders specific to foetus and newborn
P75-P78	Digestive system disorders of foetus and newborn
P80-P83	Conditions involving the integument and temperature regulation of foetus and newborn
P90-P96	Other disorders originating in the perinatal period
Q00-Q07	Congenital malformations of the nervous system
Q10-Q18	Congenital malformations of eye, ear, face and neck
Q20-Q28	Congenital malformations of the circulatory system
Q30-Q34	Congenital malformations of the respiratory system
Q35-Q37	Cleft lip and cleft palate
Q38-Q45	Other congenital malformations of the digestive system
Q50-Q56	Congenital malformations of genital organs
Q60-Q64	Congenital malformations of the urinary system
Q65-Q79	Congenital malformations and deformations of the musculoskeletal system
Q80-Q89	Other congenital malformations
Q90-Q99	Chromosomal abnormalities, not elsewhere classified

All other codes (A00 – O99, R00 - Z99) have been classified to the category 'all other causes'.

# **APPENDIX E**

#### **CLASSIFICATION OF COUNTY OF RESIDENCE**

The following codes are applied to mother's and father's county of residence. Where a Dublin postal code, such as Dublin 13, is entered then a code of 10 is applied for Dublin City. Where there is no postal code and Dublin is recorded then a code of 11 is applied for Dublin County. For births occurring in the HSE South region where Tipperary is recorded as county of residence then a code of 62, Tipperary South Riding, is applied. For births occurring in other HSE regions where Tipperary is recorded as county of residence then a code of 32, Tipperary North Riding, is applied. Where a mother's or father's county of residence is unknown the code 99 is applied. The code 90 is applied where the address given is outside the Republic of Ireland.

10	Dublin City	50	Donegal
11	Dublin County	51	Sligo
12	Kildare	52	Leitrim
13	Wicklow	60	Carlow
20	Laois	61	Kilkenny
21	Offaly	62	Tipperary South Riding
22	Longford	63	Waterford
23	Westmeath	64	Wexford
30	Clare	70	Cork
31	Limerick	71	Kerry
32	Tipperary North Riding	80	Galway
40	Meath	81	Mayo
41	Cavan	82	Roscommon
42	Monaghan	90	Other – all places outside Republic of Ireland
43	Louth	99	Not Stated
		22	NOI Stated

# **APPENDIX F**

#### **CLASSIFICATION OF NATIONALITY**

Nationality for mother and father is defined by the place of their birth. This definition is for the sole purpose of giving the necessary directions for the completion of the Birth Notification Form and is not a legal definition.

Nationalities have been coded and classified into a set of groups according to a system devised by the CSO.

#### Country groupings are as follows:

Ireland includes the following: UK includes the following:

IrelandEnglandIrish AmericanGreat BritainIrish EnglishNorthern Ireland

Irish EuropeanScotlandIrish Other.Wales.

#### EU 15 (excluding Ireland and the UK) includes the following:

Austria Italy

Belgium Luxembourg
Denmark Netherlands
Finland Portugal
France Spain
Germany Sweden.

Greece

#### EU 15 to EU 27 accession states includes the following:

Bulgaria Lithuania
Cyprus Malta
Czech Republic Poland
Estonia Slovakia
Hungary Slovenia
Latvia Romania.

#### Rest of Europe includes the following:

Albania Jersey

Andorra Liechtenstein

Armenia Macedonia, The Former Yugoslav Republic of,

Azerbaijan Moldova, Republic of

Belarus Monaco Bosnia and Herzegovina Norway

Croatia Russian Federation

Faroe Islands San Marino
Georgia Switzerland
Gibraltar Turkey
Guernsey Ukraine
Iceland Vatican City

Isle of Man Yugoslavia, Federal Republic of.

#### Africa includes the following:

Congo, The Democratic Republic of the

Malawi Algeria Angola Mali Benin Mauritania Botswana Mauritius Burkina Faso Mayotte Burundi Morocco Cameroon Mozambique Cape Verde Namibia Central African Republic Niger Chad Nigeria Comoros Reunion Rwanda Congo

Cote d'Ivoire Sao Tome and Principe

Saint Helena

Djibouti Senegal
Egypt Seychelles
Equatorial Guinea Sierra Leone
Eritrea Somalia
Ethiopia South Africa
Gabon Sudan
Gambia Swaziland

Ghana Tanzania, United Republic of

Guinea Togo Guinea-Bissau Tunisia Kenya Uganda

Lesotho Western Sahara

Liberia Zambia Libyan Arab Jamahiriya Zimbabwe.

Madagascar

#### Asia includes the following:

Afghanistan Macao
Bahrain Malaysia
Bangladesh Maldives
Bhutan Mongolia
British Indian Ocean Territory Myanmar
Brunei Darussalam Nepal

Cambodia Oman China Pakistan

East Timor Palestinian Territory, Occupied

Hong Kong Philippines
India Qatar
Indonesia Saudi Arabia

Iran, Islamic Republic ofSingaporeIraqSri Lanka

Israel Syrian Arab Republic
Japan Taiwan, Province of China

JordanTajikistanKazakhstanThailandKorea, Democratic People's Republic ofTurkmenistan

Korea, Republic of United Arab Emirates

KuwaitUzbekistanKyrgyzstanVietnamLao People's Democratic RepublicYemen.

Lebanon

#### America includes the following:

Anguilla Guyana
Antigua and Barbuda Haiti
Argentina Honduras
Aruba Jamaica
Bahamas Martinique
Barbados Mexico
Belize Montserrat

Bermuda Netherlands Antilles

Bolivia Nicaragua
Brazil Panama
Canada Paraguay
Cayman Islands Peru

Chile Puerto Rico

Colombia Saint Kitts and Nevis

Costa Rica Saint Lucia

Cuba Saint Pierre and Miquelon

Dominica Saint Vincent and the Grenadines

Dominican Republic Suriname

Ecuador Trinidad and Tobago
El Salvador Turks and Caicos Islands

Falkland Islands (Malvinas)

French Guiana

Greenland

United States

Uruguay

Venezuela

Grenada Virgin Islands, British Guadeloupe Virgin Island, US.

Guatemala

#### Australia includes the following:

Australia.

#### New Zealand and Other Oceania includes the following:

American Samoa Niue

Antarctica Norfolk Island

Bouvet Island Northern Mariana Islands

Christmas Island Palau

Cocos (Keeling) Islands Papua New Guinea

Cook Islands Pitcairn
Fiji Samoa

French Polynesia Solomon Islands

French Southern Territories South Georgia and the South Sandwich Islands

Guam Svalbard and Jan Mayen

Heard Island and McDonald Islands

Kiribati

Tonga

Marshall Islands

Tuvalu

Micronesia, Federated States of United States Minor Outlying Islands

Nauru Vanuatu

New Caledonia Wallis and Futuna.

New Zealand

#### Multi-Nationality includes the following:

Mixed/Multi-Nationality (stated as so).

#### No Nationality:

Stated as so.

#### Not Stated:

Where nationality is not provided.

#### Not Applicable:

This applies to father's nationality only. Where mother's marital status is stated as 'never married', 'widowed', 'separated' or 'divorced' the General Register Office does not require father's details to register the birth, and as such this information is generally not provided.

#### All Other Nationalities:

Where nationality is not recorded to any category listed above.

# **APPENDIX G**

## **DOMICILIARY BIRTHS, 1999-2007**

#### **INTRODUCTION**

Domiciliary births have been included, unless otherwise stated, in the analysis presented in previous sections. Domiciliary births are defined as planned home births attended by an independent midwife and are not associated with a hospital. Births delivered under hospital administered home birth schemes are recorded as hospital births.

In 1999 there were 246 domiciliary births. Of these, there was one early neonatal death and no stillbirths. In 2000 the number of domiciliary births fell to 216 with, again, one early neonatal death and no stillbirths. There were no domiciliary multiple births in 1999 or 2000.

In 2001 the number of domiciliary births increased to 245. Of these, there was one stillbirth and no early neonatal deaths. There was one set of domiciliary twin births in 2001.

The trend in the number of domiciliary births continued to be upwards, increasing to 288 in 2002. Of these, there were no stillbirths and one early neonatal death. There were no domiciliary multiple births in 2002.

In 2003 and 2004 the number of domiciliary births decreased to 236 and 202 respectively. Of these, there were no stillbirths and one early neonatal death in 2004. There were no domiciliary multiple births in 2003 or 2004. In 2005 there were 183 planned domiciliary births reported to NPRS. There were no stillbirths, early neonatal deaths or multiple births among domiciliary births in 2005.

For 2006 there were 170 domiciliary births. Again, there were no stillbirths, early neonatal deaths or multiple births among domiciliary births in 2006. There were 186 domiciliary births recorded in 2007 and there was one stillbirth and no early neonatal deaths. There were no domiciliary multiple births in 2007.

Table G1 below highlights trends across domiciliary births from 1999 to 2007 and shows that those women choosing a domiciliary delivery were older (average age 34 years), with higher maternal parity, 1.60 previous births, than those choosing to have their babies in a hospital setting (see Table 2.1, page 9). Only 16.1 per cent of mothers recorded marital status as 'never married' compared to over 30 per cent in hospital maternities. The average birthweight was higher at 3,699 grams and breastfeeding was significantly higher at over 95 per cent. Further tables show that almost 85 per cent of domiciliary births were to

multiparous women (Table G2). Over 80 per cent of domiciliary births were to women aged 30 years or more (Table G3). All but two births weighed over 2,500 grams, with over 90 per cent weighing over 3,000 grams, and all babies were born at 37 or more completed weeks gestation (Table G4). Almost two-thirds of domiciliary births were to married women aged between 30 to 39 years of age (Table G5). The most common mother's occupation was 'lower professional' followed by 'home duties' (Table G6). The highest proportion of breastfed babies where to those whose father's occupation was recorded as 'skilled manual workers', and to mother's recording an occupation of 'lower professional' (Tables G7 and G8).

The tables that follow present selected statistics on domiciliary births for 1999 to 2007 (G1) and contain a range of cross-tabulations relating to all domiciliary births for 2007 (G2-G8). The format of each cross-tabulation is identical. The values of the two variables are given along the top row and left-hand side column. The top number in each cell indicates the number of observations relating to that combination of variable values. The second number indicates the percentage contribution of that cell to the total for the whole row. The last number indicates the percentage contribution of that cell to the total for the whole column. The figures at the end of each row or column give the overall total for that row or column, and the associated percentages express each of these figures as a percentage of the grand total.

**TABLE G1: DOMICILIARY BIRTHS** 

Trends in Selected Statistics, 1999-2007

Selected Statistics	1999	2000	2001	2002	2003	2004	2005	2006	2007
General Characteristics									
Average Age of Mother in years <sup>1</sup>	33.47	33.33	33.18	34.07	34.01	34.31	33.37	34.34	34.12
Average Maternal Parity <sup>1</sup>	1.52	1.62	1.40	2.07	1.63	1.60	1.70	1.72	1.60
Single Mothers (%) 1,2	25.20	29.63	31.56	22.57	27.97	29.21	23.50	20.00	16.13
Average Weight in grams <sup>3</sup>									
Total Live Births	3,699.46	3,726.71	3,686.75	3,730.06	3,754.25	3,738.40	3,760.09	3,742.20	3,699.24
Low Birthweight <2,500 grams (%) <sup>3</sup>									
Total Live Births	0.43	1.86	0.00	0.35	0.00	0.99	0.55	0.59	1.08
Average Gestational Age at delivery in weeks	40.04	39.92	40.02	40.14	40.05	39.99	40.03	40.08	39.98
Average Interval since last birth in years	3.46	3.40	3.82	3.74	3.65	3.54	3.30	3.44	3.38
Perinatal Care									
Combined Antenatal Care - Midwife and GP/Hospital (%) <sup>4</sup>	59.76	57.87	55.33	57.29	66.10	64.36	71.58	76.47	60.75
Immunity to Rubella (%) <sup>1</sup>	92.34	90.70	93.44	91.61	94.07	97.03	95.08	96.47	100.00
Mothers Breastfeeding (%) 5	98.25	96.63	95.02	94.04	94.07	97.51	93.99	97.06	95.68

Notes: There were no domiciliary multiple births in 1999, 2000 or in the years 2002 -2007.

There was one set of domiciliary twin births in 2001.

'Not Stated' values are not included in the calculation of percentages.

<sup>&</sup>lt;sup>1</sup> Based on maternities.

<sup>&</sup>lt;sup>2</sup> Data recording marital status as 'never married' are classified as 'single' marital status in this report.

<sup>&</sup>lt;sup>3</sup> Based on live births.

<sup>&</sup>lt;sup>4</sup> Based on maternities. This figure applies to the percentage of home births where antenatal care is stated as 'combined', rather than 'midwife only'.

<sup>&</sup>lt;sup>5</sup> Based on maternities with live births. Excludes early neonatal deaths.

**TABLE G2: DOMICILIARY BIRTHS** 

Age of Mother by Maternal Parity, 2007

Age of Mother				Materna	al Parity			
Frequency	0	1	2	3	4	5	6	Total
Row Per Cent								
Col. Per Cent								
20-24 Years	1	3	0	0	0	0	0	4
	25.0	75.0	0.0	0.0	0.0	0.0	0.0	100
	3.4	4.3	0.0	0.0	0.0	0.0	0.0	2.2
25-29 Years	9	11	6	0	0	0	0	26
	34.6	42.3	23.1	0.0	0.0	0.0	0.0	100
	31.0	15.7	11.3	0.0	0.0	0.0	0.0	14.0
30-34 Year	12	27	24	10	2	1	0	76
	15.8	35.5	31.6	13.2	2.6	1.3	0.0	100
	41.4	38.6	45.3	45.5	33.3	20.0	0.0	40.9
35-39 Years	6	25	21	9	2	2	0	65
	9.2	38.5	32.3	13.8	3.1	3.1	0.0	100
	20.7	35.7	39.6	40.9	33.3	40.0	0.0	34.9
40-44 Years	1	4	2	3	2	1	1	14
	7.1	28.6	14.3	21.4	14.3	7.1	7.1	100
	3.4	5.7	3.8	13.6	33.3	20.0	100.0	7.5
45 Years and	0	0	0	0	0	1	0	1
Over	0.0	0.0	0.0	0.0	0.0	100.0	0.0	100
	0.0	0.0	0.0	0.0	0.0	20.0	0.0	0.5
Total	29	70	53	22	6	5	1	186
	15.6	37.6	28.5	11.8	3.2	2.7	0.5	100
	100	100	100	100	100	100	100	100

**TABLE G3: DOMICILIARY BIRTHS** 

Age of Mother by Gestational Age at Delivery, 2007

Age of Mother		Gestational Age (weeks	)
Frequency	37-41	42 and Over	Total
Row Per Cent			
Col. Per Cent			
20-24 Years	3	1	4
	75.0	25.0	100
	1.7	12.5	2.2
25-29 Years	26	0	26
	100.0	0.0	100
	14.6	0.0	14.0
30-34 Year	72	4	76
	94.7	5.3	100
	40.4	50.0	40.9
35-39 Years	62	3	65
	95.4	4.6	100
	34.8	37.5	34.9
40-44 Years	14	0	14
	100.0	0.0	100
	7.9	0.0	7.5
45 Years and Over	1	0	1
	100.0	0.0	100
	0.6	0.0	0.5
Total	178	8_	186
	95.7	4.3	100
	100	100	100

**TABLE G4: DOMICILIARY BIRTHS** 

Birthweight by Gestational Age at Delivery, 2007

Birthweight (grams)	Gestational Age (weeks)						
Frequency	37-41	42 and Over	Total				
Row Per Cent							
Col. Per Cent							
2000-2499	2	0	2				
	100.0	0.0	100				
2700 2000	1.1	0.0	1.1				
2500-2999	10	0	10				
	100.0	0.0	100				
2000 2400	5.6	0.0	5.4				
3000-3499	47	1	48				
	97.9	2.1	100				
3500 3000	26.4	12.5	25.8				
3500-3999	74	3	77				
	96.1	3.9	100				
4000 4400	41.6	37.5	41.4				
4000-4499	28	3	31				
	90.3	9.7	100				
4500 10	15.7	37.5	16.7				
4500 and Over	17	1	18				
	94.4	5.6	100				
	9.6	12.5	9.7				
Total	178	8_	186				
	95.7	4.3	100				
	100	100	100				

**TABLE G5: DOMICILIARY BIRTHS** 

Mother's Marital Status by Age of Mother, 2007

Mother's Marital Status	Age of Mother								
Frequency	20-24	25-29	30-34	35-39	40-44	45 Years	Total		
Row Per Cent	Years	Years	Years	Years	Years	and Over			
Col. Per Cent									
Married	3	15	61	55	12	1	147		
	2.0	10.2	41.5	37.4	8.2	0.7	100		
	75.0	57.7	80.3	84.6	85.7	100.0	79.0		
Never Married	1	10	11	7	1	0	30		
	3.3	33.3	36.7	23.3	3.3	0.0	100		
	25.0	38.5	14.5	10.8	7.1	0.0	16.1		
Separated	0	1	3	2	1	0	7		
	0.0	14.3	42.9	28.6	14.3	0.0	100		
	0.0	3.8	3.9	3.1	7.1	0.0	3.8		
Divorced	0	0	1	1	0	0	2		
	0.0	0.0	50.0	50.0	0.0	0.0	100		
	0.0	0.0	1.3	1.5	0.0	0.0	1.1		
Total	4	26	76	65	14	1	186		
	2.2	14.0	40.9	34.9	7.5	0.5	100		
	100	100	100	100	100	100	100		

# Mother's Occupation by Birthweight, 2007

Mother's Occupation			Birt	hweight (gra	ms)		
Frequency	2000-	2500-	3000-	3500-	4000-	4500 and	Total
Row Per Cent	2499	2999	3499	3999	4499	Over	
Col. Per Cent							
Farmers & Farm	0	0	0	0	0	1	1
Managers	0.0	0.0	0.0	0.0	0.0	100.0	100
	0.0	0.0	0.0	0.0	0.0	5.6	0.5
Higher Professional	1	1	2	8	4	4	20
	5.0	5.0	10.0	40.0	20.0	20.0	100
	50.0	10.0	4.2	10.4	12.9	22.2	10.8
Lower Professional	0	4	15	27	7	3	56
	0.0	7.1	26.8	48.2	12.5	5.4	100
	0.0	40.0	31.3	35.1	22.6	16.7	30.1
Employers &	0	0	1	3	0	0	4
Managers	0.0	0.0	25.0	75.0	0.0	0.0	100
	0.0	0.0	2.1	3.9	0.0	0.0	2.2
Salaried Employees	0	0	1	0	0	0	1
	0.0	0.0	100.0	0.0	0.0	0.0	100
	0.0	0.0	2.1	0.0	0.0	0.0	0.5
Intermediate Non- Manual Workers	0	0	6	8	3	2	19
Manual Workers	0.0	0.0	31.6	42.1	15.8	10.5	100
	0.0	0.0	12.5	10.4	9.7	11.1	10.2
Other Non-Manual Workers	0	1	5	10	6	3	25
WOIKEIS	0.0	4.0	20.0	40.0	24.0	12.0	100
Chille d Manual	0.0	10.0	10.4	13.0	19.4	16.7	13.4
Skilled Manual Workers	0	0	1	0	2	0	3
WOIKEIS	0.0	0.0	33.3	0.0	66.7	0.0	100
Not Classifiable	0.0	0.0	2.1	0.0	6.5	0.0	1.6
NOT Classifiable	0	2	0	3	2	1	8
	0.0	25.0	0.0	37.5	25.0	12.5	100
Home Duties	0.0	20.0	0.0	3.9	6.5	5.6	4.3
Home Duties	1	2	17	18	7	4	49
	2.0	4.1	34.7	36.7	14.3	8.2	100
Total	50.0	20.0	35.4	23.4	22.6	22.2	26.3
Total	2	10	48	77	31	18	186
	1.1	5.4	25.8	41.4	16.7	9.7	100
	100	100	100	100	100	100	100

Father's Occupation by Infant's Type of Feeding, Live Births, 2007

Father's Occupation	Type of Feeding					
Frequency	Artificial	Breast	Combined	Total		
Row Per Cent		_		_		
Col. Per Cent		_		_		
Farmers & Farm Managers	0	4	0	4		
	0.0 0.0	100.0 2.3	0.0 0.0	100 2.2		
Other Agricultural Occupations	0.0	1	0.0	1		
& Fishermen	0.0	100.0	0.0	100		
	0.0	0.6	0.0	0.5		
Higher Professional	0	22	0	22		
	0.0	100.0	0.0	100		
	0.0	12.4	0.0	11.9		
Lower Professional	1	24	0	25		
	4.0	96.0	0.0	100		
	14.3	13.6	0.0	13.5		
Employers & Managers	2	21	0	23		
	8.7	91.3	0.0	100		
	28.6	11.9	0.0	12.4		
Salaried Employees	0	3	0	3		
	0.0	100.0	0.0	100		
Internal distable New Manage	0.0	1.7	0.0	1.6		
Intermediate Non-Manual Workers	0	17	0	17		
WORKERS	0.0	100.0	0.0	100		
Other Non-Manual Workers	0.0	9.6	0.0	9.2		
Other Non-Ivialidal Workers	0	13	0	13		
	0.0 0.0	100.0 7.3	0.0 0.0	100 7.0		
Skilled Manual Workers	0.0	7.3 58	0.0	61		
Skilled Warrach Workers	3.3	95.1	1.6	100		
	28.6	32.8	100.0	33.0		
Semi-Skilled Manual Workers	0	1	0	1		
	0.0	100.0	0.0	100		
	0.0	0.6	0.0	0.5		
Unskilled Manual Workers	0	1	0	1		
	0.0	100.0	0.0	100		
	0.0	0.6	0.0	0.5		
Unemployed	0	2	0	2		
	0.0	100.0	0.0	100		
	0.0	1.1	0.0	1.1		
Not Classifiable	2	8	0	10		
	20.0	80.0	0.0	100		
	28.6	4.5	0.0	5.4		
Not Applicable	0	2	0	2		
	0.0	100.0	0.0	100		
Tabel	0.0	1.1	0.0	1.1		
Total	7_	177_	11	185		
	3.8	95.7	0.5	100		
	100	100	100	100		

**TABLE G8: DOMICILIARY BIRTHS** 

Mother's Occupation by Infant's Type of Feeding, Live Births, 2007

Mother's Occupation	Type of Feeding							
Frequency	Artificial	Breast	Combined	Total				
Row Per Cent								
Col. Per Cent								
Farmers & Farm Managers	0	1	0	1				
	0.0	100.0	0.0	100				
	0.0	0.6	0.0	0.5				
Higher Professional	1	19	0	20				
	5.0	95.0	0.0	100				
	14.3	10.7	0.0	10.8				
Lower Professional	1	53	1	55				
	1.8	96.4	1.8	100				
	14.3	29.9	100.0	29.7				
Employers & Managers	0	4	0	4				
	0.0	100.0	0.0	100				
	0.0	2.3	0.0	2.2				
Salaried Employees	0	1	0	1				
	0.0	100.0	0.0	100				
Internaciate New Manual	0.0	0.6	0.0	0.5				
Intermediate Non-Manual Workers	3	16	0	19				
Workers	15.8	84.2	0.0	100				
Other Non-Manual Workers	42.9	9.0	0.0	10.3				
Other Non-Ivianual Workers	0	25	0	25				
	0.0	100.0 14.1	0.0	100				
Skilled Manual Workers	0.0	14.1	0.0	13.5 3				
Skilled Walladi Workers	0.0	100.0	0.0	100				
	0.0	1.7	0.0	1.6				
Not Classifiable	0.0	7	0.0	1.6				
	12.5	87.5	0.0	100				
	14.3	4.0	0.0	4.3				
Home Duties	14.5	48	0.0	49				
	2.0	98.0	0.0	100				
	14.3	27.1	0.0	26.5				
Total	7	177	1	185				
	3.8	95.7	0.5	100				
	100	100	100	100				

# **APPENDIX H**

# **SOURCE DATA FOR FIGURES**

**TABLE H1: TOTAL BIRTHS** 

Age of Mother

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Age of Mother	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
< 20 Years	2,507	3.5	14	5.6	8	3.2	8.7	6.0
20-24 Years	8,939	12.5	55	6.1	18	2.0	8.1	6.7
25-29 Years	16,617	23.2	83	5.0	22	1.3	6.3	4.9
30-34 Years	24,493	34.2	105	4.3	48	2.0	6.2	4.3
35+ Years	18,998	26.5	130	6.8	36	1.9	8.7	6.1
Not Stated	16	0.0	6	272.7	0	0.0	272.7	272.7
Total	71,570	100	393	5.5	132	1.8	7.3	5.3

**TABLE H2: TOTAL BIRTHS** 

Percentage of Births by Mother's Marital Status, 2003-2007

Marital Status	2003	2004	2005	2006	2007
Married	68.4	67.7	67.6	66.8	66.4
Never Married	30.1	30.5	30.5	31.4	31.7
Widowed	0.1	0.1	0.1	0.2	0.1
Separated	1.1	1.2	1.2	1.1	1.1
Divorced	0.3	0.4	0.5	0.6	0.6

**TABLE H3: TOTAL BIRTHS**Mother's County of Residence by Mother's Marital Status, 2007

Mother's County of Residence	Mother's Marital Status								
Frequency	Married	Never	Widowed	Separated	Divorced	Not	Total		
Row Per Cent		Married				Stated			
Col. Per Cent	620	252	1				006		
Carlow	629 63.8	352 35.7	1 0.1	4 0.4	0.0	0 0.0	986 100		
	1.3	1.5	1.1	0.4	0.0	0.0	1.4		
Cavan	782	291	3	13	12	0	1,101		
	71.0	26.4	0.3	1.2	1.1	0.0	100		
	1.6	1.3	3.4	1.6	2.8	0.0	1.5		
Clare	1,300	482	3	15	7	0	1,807		
	71.9 2.7	26.7 2.1	0.2 3.4	0.8 1.8	0.4 1.7	0.0 0.0	100 2.5		
Cork	5,624	2,210	16	88	59	0.0	7,997		
<b>3</b> 0	70.3	27.6	0.2	1.1	0.7	0.0	100		
	11.8	9.7	18.0	10.6	14.0	0.0	11.1		
Donegal	1,452	638	2	30	22	0	2,144		
	67.7	29.8	0.1	1.4	1.0	0.0	100		
Dublin City	3.0	2.8	2.2	3.6	5.2	0.0	3.0		
Dublin City	8,421 57.7	5,909 40.5	18 0.1	186 1.3	72 0.5	1 0.0	14,607 100		
	17.6	25.9	20.2	22.5	17.1	16.7	20.3		
Dublin County	3,958	1,515	5	58	27	0	5,563		
	71.1	27.2	0.1	1.0	0.5	0.0	100		
	8.3	6.6	5.6	7.0	6.4	0.0	7.7		
Galway	2,838	931	2	47	24	0	3,842		
	73.9 5.9	24.2 4.1	0.1 2.2	1.2 5.7	0.6 5.7	0.0 0.0	100 5.3		
Kerry	1,471	551	2.2	28	11	0.0	2,063		
Refry	71.3	26.7	0.1	1.4	0.5	0.0	100		
	3.1	2.4	2.2	3.4	2.6	0.0	2.9		
Kildare	2,786	1,086	1	33	12	0	3,918		
	71.1	27.7	0.0	0.8	0.3	0.0	100		
William on the	5.8	4.8	1.1	4.0	2.8	0.0	5.4		
Kilkenny	928 72.3	348 27.1	0.0	5 0.4	0.2	0 0.0	1,283 100		
	1.9	1.5	0.0	0.4	0.2	0.0	1.8		
Laois	866	400	2	14	3	1	1,286		
	67.3	31.1	0.2	1.1	0.2	0.1	100		
	1.8	1.8	2.2	1.7	0.7	16.7	1.8		
Leitrim	347	130	2	7	2	0	488		
	71.1	26.6	0.4	1.4	0.4	0.0	100		
Limerick	0.7 1,916	0.6 1,053	2.2	0.8 28	0.5 13	0.0	0.7 3,016		
Limerick	63.5	34.9	0.2	0.9	0.4	0.0	100		
	4.0	4.6	6.7	3.4	3.1	0.0	4.2		
Longford	410	168	1	16	5	0	600		
	68.3	28.0	0.2	2.7	0.8	0.0	100		
	0.9	0.7	1.1	1.9	1.2	0.0	0.8		

TABLE H3: TOTAL BIRTHS (contd.)

Mother's County of Residence by Mother's Marital Status, 2007

Mother's County of Residence	Mother's Marital Status							
Frequency	Married	Never	Widowed	Separated	Divorced	Not	Total	
Row Per Cent		Married				Stated	_	
Col. Per Cent	_				_	_	_	
Louth	1,088	712	3	28	13	0	1,844	
	59.0	38.6	0.2	1.5	0.7	0.0	100	
	2.3	3.1	3.4	3.4	3.1	0.0	2.6	
Mayo	1,353	462	4	22	21	2	1,864	
,	72.6	24.8	0.2	1.2	1.1	0.1	100	
	2.8	2.0	4.5	2.7	5.0	33.3	2.6	
Meath	2,478	974	0	26	32	0	3,510	
	70.6	27.7	0.0	0.7	0.9	0.0	100	
	5.2	4.3	0.0	3.1	7.6	0.0	4.9	
Monaghan	570	230	0	10	5	0	815	
	69.9	28.2	0.0	1.2	0.6	0.0	100	
	1.2	1.0	0.0	1.2	1.2	0.0	1.1	
Offaly	771	388	0	9	4	0	1,172	
	65.8	33.1	0.0	0.8	0.3	0.0	100	
	1.6	1.7	0.0	1.1	0.9	0.0	1.6	
Roscommon	562	185	0	13	2	0	762	
	73.8	24.3	0.0	1.7	0.3	0.0	100	
	1.2	0.8	0.0	1.6	0.5	0.0	1.1	
Sligo	616	279	0	12	7	0	914	
	67.4	30.5	0.0	1.3	0.8	0.0	100	
	1.3	1.2	0.0	1.5	1.7	0.0	1.3	
Tipperary North	651	257	3	12	2	0	925	
	70.4	27.8	0.3	1.3	0.2	0.0	100	
	1.4	1.1	3.4	1.5	0.5	0.0	1.3	
Tipperary South	917	536	4	14	7	0	1,478	
	62.0	36.3	0.3	0.9	0.5	0.0	100	
	1.9	2.3	4.5	1.7	1.7	0.0	2.1	
Waterford	1,144	661	0	21	14	0	1,840	
	62.2	35.9	0.0	1.1	0.8	0.0	100	
	2.4	2.9	0.0	2.5	3.3	0.0	2.6	
Westmeath	955	429	3	20	10	1	1,418	
	67.3	30.3	0.2	1.4	0.7	0.1	100	
	2.0	1.9	3.4	2.4	2.4	16.7	2.0	
Wexford	1,359	872	3	40	25	0	2,299	
	59.1	37.9	0.1	1.7	1.1	0.0	100	
	2.8	3.8	3.4	4.8	5.9	0.0	3.2	
Wicklow	1,600	764	5	28	7	1	2,405	
	66.5	31.8	0.2	1.2	0.3	0.0	100	
	3.3	3.3	5.6	3.4	1.7	16.7	3.3	
Other	8	6	0	0	2	0	16	
	50.0	37.5	0.0	0.0	12.5	0.0	100	
	0.0	0.0	0.0	0.0	0.5	0.0	0.0	
Total	47,800	22,819	89	827	422	6	71,963	
	66.4	31.7	0.1	1.1	0.6	0.0	100	
	100	100	100	100	100	100	100	

*Note*: See Appendix E for the classification system employed for county of residence.

**TABLE H4: TOTAL BIRTHS** 

Mother's Occupation

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Mother's Occupation	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Farmers & Farm Managers	65	0.1	2	29.9	0	0.0	29.9	29.9
Other Agricultural Occupations & Fishermen	93	0.1	0	0.0	0	0.0	0.0	0.0
Higher Professional	4,323	6.0	23	5.3	7	1.6	6.9	4.2
Lower Professional	9,905	13.8	43	4.3	17	1.7	6.0	4.2
Employers & Managers	5,121	7.2	22	4.3	13	2.5	6.8	4.3
Salaried Employees	1,085	1.5	7	6.4	3	2.8	9.2	5.5
Intermediate Non-Manual Workers	16,371	22.9	62	3.8	23	1.4	5.2	3.8
Other Non- Manual Workers	8,904	12.4	42	4.7	9	1.0	5.7	4.4
Skilled Manual Workers	1,283	1.8	5	3.9	3	2.3	6.2	5.4
Semi-Skilled Manual Workers	2,448	3.4	16	6.5	4	1.6	8.1	6.1
Unskilled Manual Workers	7	0.0	0	0.0	0	0.0	0.0	0.0
Unemployed	2,354	3.3	27	11.3	4	1.7	13.0	10.5
Not Classifiable	2,666	3.7	23	8.6	8	3.0	11.5	9.7
Home Duties	16,795	23.5	116	6.9	40	2.4	9.2	6.8
Not Stated	150	0.2	5	32.3	1	6.7	38.7	32.5
Total	71,570	100	393	5.5	132	1.8	7.3	5.3

**TABLE H5: TOTAL BIRTHS** 

Father's Occupation

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Father's Occupation	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Farmers & Farm Managers	2,163	3.0	9	4.1	5	2.3	6.4	4.2
Other Agricultural Occupations & Fishermen	728	1.0	4	5.5	0	0.0	5.5	5.5
Higher Professional	5,507	7.7	25	4.5	6	1.1	5.6	3.4
Lower Professional	3,387	4.7	17	5.0	5	1.5	6.5	4.7
Employers & Managers	5,932	8.3	23	3.9	10	1.7	5.5	3.7
Salaried Employees	1,578	2.2	5	3.2	1	0.6	3.8	2.5
Intermediate Non-Manual Workers	5,693	8.0	29	5.1	9	1.6	6.6	5.1
Other Non- Manual Workers	6,809	9.5	32	4.7	9	1.3	6.0	4.2
Skilled Manual Workers	17,933	25.1	64	3.6	31	1.7	5.3	3.8
Semi-Skilled Manual Workers	3,211	4.5	15	4.6	8	2.5	7.1	4.4
Unskilled Manual Workers	843	1.2	4	4.7	2	2.4	7.1	2.4
Unemployed	2,176	3.0	17	7.8	5	2.3	10.0	6.4
Not Classifiable	1,582	2.2	8	5.0	2	1.3	6.3	5.7
Not Applicable	13,542	18.9	123	9.0	36	2.7	11.6	9.5
Home Duties	76	0.1	1	13.0	0	0.0	13.0	0.0
Not Stated	410	0.6	17	39.8	3	7.3	46.8	33.3
Total	71,570	100	393	5.5	132	1.8	7.3	5.3

**TABLE H6: TOTAL BIRTHS** 

Percentage of Births by Mother's Nationality, 2004-2007

Mother's Nationality	2004	2005	2006	2007
Ireland	84.4	83.5	81.6	79.3
UK	3.1	3.4	3.3	3.0
EU15 to EU27 Accession States	0.9	1.9	3.9	7.2
Africa	5.6	4.4	4.0	3.6
Asia	2.5	2.9	3.1	3.6

Notes:

Percentages calculated for the five most frequent nationalities in 2007. 'Not Stated' values are not included in the calculation of percentages.

On 1 January 2007, Bulgaria and Romania joined the European Union, giving 27 Member States. Prior to 2007 there were 25 Member States in the European Union. For 2007, less than 1 per cent of total births were to mothers whose nationality is recorded as either Bulgarian or Romanian.

**TABLE H7: TOTAL BIRTHS** 

Mother's Average Age by Mother's Nationality, 2007

Mother's Nationality	Average Age in Years
Ireland	31.5
UK	32.0
EU15 (excluding Ireland & UK)	32.5
EU15 to EU27 Accession States	27.0
Rest of Europe	29.3
Africa	30.6
Asia	30.0
America	31.3
Australia	32.4
Total	31.1

Note:

Average ages calculated for nationalities with > 100 births.

#### **TABLE H8: TOTAL BIRTHS**

Percentage of Births to Primiparous Women by Maternal Age, 2003 and 2007

Maternal Age (years)	2003	2007
<20 Years	10.2	7.6
20-24 Years	20.6	19.6
25-29 Years	27.4	27.2
30-34 Years	29.6	31.0
35-39 Years	10.3	12.4
40+ Years	1.7	2.1
Not Stated	0.1	0.0
Total	100	100

**TABLE H9: TOTAL BIRTHS** Percentage of Births to Multiparous Women by Maternal Age, 2003 and 2007

Maternal Age (years)	2003	2007
<20 Years	0.8	0.5
20-24 Years	9.0	7.3
25-29 Years	21.5	20.3
30-34 Years	37.3	36.5
35-39 Years	26.1	29.3
40+ Years	5.2	6.0
Not Stated	0.1	0.0
Total	100	100

**TABLE H10: TOTAL BIRTHS** Age of Mother by Maternal Parity, 2007

Age of Mother					Materna	al Parity				
Frequency	0	1	2	3	4	5	6	More	Not	Total
Row Per Cent								than 6	Stated	
< 20 Years	2,302	197	19	1	0	1	0	0	1	2,521
	91.3	7.8	0.8	0.0	0.0	0.0	0.0	0.0	0.0	100
20-24 Years	5,925	2,281	624	123	33	6	0	0	2	8,994
	65.9	25.4	6.9	1.4	0.4	0.1	0.0	0.0	0.0	100
25-29 Years	8,195	5,391	2,154	661	197	68	23	8	3	16,700
	49.1	32.3	12.9	4.0	1.2	0.4	0.1	0.0	0.0	100
30-34 Years	9,346	8,894	4,177	1,424	434	189	70	58	6	24,598
	38.0	36.2	17.0	5.8	1.8	0.8	0.3	0.2	0.0	100
35-39 Years	3,735	5,321	4,218	1,788	569	196	85	78	5	15,995
	23.4	33.3	26.4	11.2	3.6	1.2	0.5	0.5	0.0	100
40+ Years	643	847	748	469	222	99	55	49	1	3,133
	20.5	27.0	23.9	15.0	7.1	3.2	1.8	1.6	0.0	100
Not Stated	10	7	2	2	0	0	0	1	0	22
	45.5	31.8	9.1	9.1	0.0	0.0	0.0	4.5	0.0	100
Total	30,156	22,938	11,942	4,468	1,455	559	233	194	18	71,963
	41.9	31.9	16.6	6.2	2.0	0.8	0.3	0.3	0.0	100

#### **TABLE H11: TOTAL LIVE BIRTHS**

Birthweight Groups, Numbers and Percentages, 2007

Birthweight Groups	Number	Per Cent
Less than 1,500 grams	639	0.9
1,500 - 2,499 grams	3,127	4.4
2,500 - 4,499 grams	65,825	92.0
4,500 grams and over	1,969	2.8
Not Stated	10	0.0
Total	71,570	100

**TABLE H12: TOTAL LIVE BIRTHS** 

Birthweight Groups by Sex, 2007

Birthweight Group	Ma	ale	Female		Unkı	nown	Total	
	Number	Per Cent						
500-749	48	0.1	57	0.2	0	0.0	105	0.1
750-999	66	0.2	66	0.2	0	0.0	132	0.2
1000-1249	93	0.3	86	0.2	0	0.0	179	0.3
1250-1499	103	0.3	119	0.3	1	25.0	223	0.3
1500-1999	405	1.1	412	1.2	0	0.0	817	1.1
2000-2499	1,050	2.9	1,260	3.6	0	0.0	2,310	3.2
2500-2999	3,576	9.8	4,650	13.3	2	50.0	8,228	11.5
3000-3499	10,874	29.7	12,557	35.9	0	0.0	23,431	32.7
3500-3999	13,102	35.8	11,404	32.6	0	0.0	24,506	34.2
4000-4499	5,900	16.1	3,760	10.8	0	0.0	9,660	13.5
4500 and Over	1,382	3.8	587	1.7	0	0.0	1,969	2.8
Not Stated	8	0.0	1	0.0	1	25.0	10	0.0
Total	36,607	100	34,959	100	4	100	71,570	100

#### **TABLE H13: TOTAL BIRTHS**

Gestational Age at Delivery

Live Births, Stillbirths, Early Neonatal Deaths, and Mortality Rates, 2007

Gestational Age (weeks)	Live Births	Per Cent (%) Live Births	Stillbirths	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Less than 22	1	0.0	7	875.0	1	1,000.0	1,000.0	1,000.0
22-27	212	0.3	72	253.5	53	250.0	440.1	417.6
28-31	515	0.7	65	112.1	16	31.1	139.7	104.1
32-36	3,422	4.8	91	25.9	41	12.0	37.6	20.0
37-41	64,951	90.8	155	2.4	21	0.3	2.7	2.0
42 and Over	2,459	3.4	3	1.2	0	0.0	1.2	0.8
Not Stated	10	0.0	0	0.0	0	0.0	0.0	0.0
Total	71,570	100	393	5.5	132	1.8	7.3	5.3

**TABLE H14: SINGLETON BIRTHS**Method of Delivery by Mother's Postnatal Length of Stay, 2007

Method of Delivery			Postnatal Ler	ngth of Stay		
Frequency	0-2 Days	3-5 Days	6-8 Days	9+ Days	Not Stated	Total
Row Per Cent						
Col. Per Cent						
Spontaneous	27,795	12,401	302	59	224	40,781
	68.2	30.4	0.7	0.1	0.5	100
	83.3	36.6	15.8	20.7	98.7	58.5
Breech / Forceps	144	141	10	1	0	296
	48.6	47.6	3.4	0.3	0.0	100
	0.4	0.4	0.5	0.4	0.0	0.4
Forceps	968	1,555	59	10	2	2,594
	37.3	59.9	2.3	0.4	0.1	100
	2.9	4.6	3.1	3.5	0.9	3.7
Vacuum extraction	4,091	4,267	152	19	0	8,529
	48.0	50.0	1.8	0.2	0.0	100
	12.3	12.6	7.9	6.7	0.0	12.2
Caesarean section	344	15,452	1,384	195	1	17,376
	2.0	88.9	8.0	1.1	0.0	100
	1.0	45.6	72.3	68.4	0.4	24.9
Other	32	71	7	1	0	111
	28.8	64.0	6.3	0.9	0.0	100
	0.1	0.2	0.4	0.4	0.0	0.2
Not stated	2	3	0	0	0	5
	40.0	60.0	0.0	0.0	0.0	100
	0.0	0.0	0.0	0.0	0.0	0.0
Total	33,376	33,890	1,914	285	227	69,692
	47.9	48.6	2.7	0.4	0.3	100
	100	100	100	100	100	100

**TABLE H15: MULTIPLE BIRTHS** 

Method of Delivery by Mother's Postnatal Length of Stay, 2007

Method of Delivery		Post	natal Length of S	tay	
Frequency	0-2 Days	3-5 Days	6-8 Days	9+ Days	Total
Row Per Cent					
Col. Per Cent					
Spontaneous	118	342	35	3	498
	23.7	68.7	7.0	0.6	100
	59.9	20.9	9.5	4.5	21.9
Breech / Forceps	21	70	10	3	104
	20.2	67.3	9.6	2.9	100
	10.7	4.3	2.7	4.5	4.6
Forceps	9	47	6	0	62
	14.5	75.8	9.7	0.0	100
	4.6	2.9	1.6	0.0	2.7
Vacuum extraction	31	100	18	2	151
	20.5	66.2	11.9	1.3	100
	15.7	6.1	4.9	3.0	6.6
Caesarean section	17	1,070	298	59	1,444
	1.2	74.1	20.6	4.1	100
	8.6	65.4	80.5	88.1	63.6
Other	1	7	3	0	11
	9.1	63.6	27.3	0.0	100
	0.5	0.4	0.8	0.0	0.5
Not stated	0	1	0	0	1
	0.0	100.0	0.0	0.0	100
	0.0	0.1	0.0	0.0	0.0
Total	197	1,637	370	67	2,271
	8.7	72.1	16.3	3.0	100
	100	100	100	100	100

**TABLE H16: TOTAL LIVE BIRTHS** 

Age of Mother by Infant's Type of Feeding, 2007

Age of Mother	Type of Feeding								
Frequency	Artificial	Breast	Combined	Not Stated	Total				
Row Per Cent		_	_						
<20 Years	1,834	575	91	7	2,507				
	73.2	22.9	3.6	0.3	100				
20-24 Years	5,196	3,280	443	20	8,939				
	58.1	36.7	5.0	0.2	100				
25-29 Years	8,334	7,295	970	18	16,617				
	50.2	43.9	5.8	0.1	100				
30-34 Years	11,391	11,686	1,381	35	24,493				
	46.5	47.7	5.6	0.1	100				
35-39 Years	7,238	7,746	881	21	15,886				
	45.6	48.8	5.5	0.1	100				
40-44 Years	1,307	1,508	180	3	2,998				
	43.6	50.3	6.0	0.1	100				
45 Years and Over	50	58	5	1	114				
	43.9	50.9	4.4	0.9	100				
Not Stated	12	4	0	0	16				
	75.0	25.0	0.0	0.0	100				
Total	35,362	32,152	3,951	105	71,570				
	49.4	44.9	5.5	0.1	100				

**TABLE H17: TOTAL LIVE BIRTHS** Mother's County of Residence by Infant's Type of Feeding, 2007

Mother's County of Residence	Type of Feeding								
Frequency	Artificial	Breast	Combined	Not Stated	Total				
Row Per Cent				_					
Col. Per Cent									
Carlow	514	454	12	0	980				
	52.4	46.3	1.2	0.0	100				
Carra	1.5	1.4	0.3	0.0	1.4				
Cavan	589 53.8	472 43.1	32 2.9	1 0.1	1,094 100				
	1.7	1.5	0.8	1.0	1.5				
Clare	1,303	420	71	5	1,799				
Ciai C	72.4	23.3	3.9	0.3	100				
	3.7	1.3	1.8	4.8	2.5				
Cork	3,847	3,608	495	10	7,960				
	48.3	45.3	6.2	0.1	100				
	10.9	11.2	12.5	9.5	11.1				
Donegal	1,302	666	156	5	2,129				
	61.2	31.3	7.3	0.2	100				
	3.7	2.1	3.9	4.8	3.0				
Dublin City	6,499	7,008	987	24	14,518				
	44.8	48.3	6.8	0.2	100				
Dublin County	18.4	21.8	25.0	22.9	20.3				
Dubilii County	2,104 38.1	3,025 54.7	394 7.1	6 0.1	5,529 100				
	5.9	9.4	10.0	5.7	7.7				
Galway	1,688	1,577	560	1	3,826				
<i>,</i>	44.1	41.2	14.6	0.0	100				
	4.8	4.9	14.2	1.0	5.3				
Kerry	1,061	890	94	5	2,050				
	51.8	43.4	4.6	0.2	100				
	3.0	2.8	2.4	4.8	2.9				
Kildare	1,742	1,966	183	5	3,896				
	44.7	50.5	4.7	0.1	100				
rell	4.9	6.1	4.6	4.8	5.4				
Kilkenny	595	678	5	2	1,280				
	46.5	53.0 2.1	0.4 0.1	0.2	100 1.8				
Laois	1.7 628	615	35	1.9 1	1,279				
Luois	49.1	48.1	2.7	0.1	100				
	1.8	1.9	0.9	1.0	1.8				
Leitrim	243	211	30	1	485				
	50.1	43.5	6.2	0.2	100				
	0.7	0.7	0.8	1.0	0.7				
Limerick	2,289	645	56	6	2,996				
	76.4	21.5	1.9	0.2	100				
	6.5	2.0	1.4	5.7	4.2				
Longford	285	277	26	3	591				
	48.2	46.9	4.4	0.5	100				
	0.8	0.9	0.7	2.9	0.8				

**TABLE H17: TOTAL LIVE BIRTHS (contd.)** 

Mother's County of Residence by Infant's Type of Feeding, 2007

Mother's County of Residence		T	ype of Feeding		
Frequency	Artificial	Breast	Combined	Not Stated	Total
Row Per Cent			_	_	
Col. Per Cent					
Louth	980	758	94	1	1,833
	53.5	41.4	5.1	0.1	100
	2.8	2.4	2.4	1.0	2.6
Mayo	828	954	77	1	1,860
	44.5	51.3	4.1	0.1	100
	2.3	3.0	1.9	1.0	2.6
Meath	1,596	1,765	131	3	3,495
	45.7	50.5	3.7	0.1	100
	4.5	5.5	3.3	2.9	4.9
Monaghan	437	356	21	0	814
	53.7	43.7	2.6	0.0	100
Official .	1.2	1.1	0.5	0.0	1.1
Offaly	617 52.9	531	15 1.3	3	1,166 100
	1.7	45.5 1.7	0.4	0.3 2.9	1.6
Roscommon	380	344	35	2.9	
ROSCOTITION	50.1	45.3	4.6	0.0	759 100
	1.1	1.1	0.9	0.0	1.1
Sligo	479	331	95	2	907
31150	52.8	36.5	10.5	0.2	100
	1.4	1.0	2.4	1.9	1.3
Tipperary North	633	277	9	1.5	920
,	68.8	30.1	1.0	0.1	100
	1.8	0.9	0.2	1.0	1.3
Tipperary South	823	634	11	3	1,471
	55.9	43.1	0.7	0.2	100
	2.3	2.0	0.3	2.9	2.1
Waterford	913	882	33	1	1,829
	49.9	48.2	1.8	0.1	100
	2.6	2.7	0.8	1.0	2.6
Westmeath	644	737	29	3	1,413
	45.6	52.2	2.1	0.2	100
	1.8	2.3	0.7	2.9	2.0
Wexford	1,244	921	112	6	2,283
	54.5	40.3	4.9	0.3	100
	3.5	2.9	2.8	5.7	3.2
Wicklow	1,094	1,140	153	5	2,392
	45.7	47.7	6.4	0.2	100
Other	3.1	3.5	3.9	4.8	3.3
Other	5	10	0	1	16
	31.3	62.5	0.0	6.3	100
Total	0.0	0.0	0.0	1.0	0.0
Total	35,362	32,152	3,951	105	71,570
	49.4	44.9	5.5	0.1	100
	100	100	100	100	100

*Note*: See Appendix E for the classification system employed for county of residence.

**TABLE H18: TOTAL BIRTHS** Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal Deaths, and Mortality Rates by Birthweight, 2007

Birthweight (grams)	Live Births	Per Cent (%) Live Births	Ante- partum Stillbirths	Intra- partum Stillbirths	Not Stated	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
500-749	105	0.1	58	6	4	393.1	42	400.0	635.8	615.9
750-999	132	0.2	28	1	2	190.2	17	128.8	294.5	243.4
1000-1249	179	0.3	30	0	0	143.5	6	33.5	172.2	121.8
1250-1499	223	0.3	23	2	1	104.4	9	40.4	140.6	77.6
1500-1999	817	1.1	34	4	5	50.0	16	19.6	68.6	39.6
2000-2499	2,310	3.2	45	2	1	20.4	18	7.8	28.0	14.6
2500-2999	8,228	11.5	45	3	1	5.9	10	1.2	7.1	5.4
3000-3499	23,431	32.7	43	4	5	2.2	7	0.3	2.5	2.2
3500-3999	24,506	34.2	21	4	3	1.1	5	0.2	1.3	1.0
4000-4499	9,660	13.5	7	1	2	1.0	2	0.2	1.2	0.9
4500 and Over	1,969	2.8	2	0	0	1.0	0	0.0	1.0	1.0
Not Stated	10	0.0	6	0	0	375.0	0	0.0	375.0	285.7
Total	71,570	100	342	27	24	5.5	132	1.8	7.3	5.3

#### **TABLE H19: TOTAL BIRTHS**

Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal Deaths, and Mortality Rates by Gestational Age, 2007

Gestational Age at Delivery (weeks)	Live Births	Per Cent (%) Live Births	Ante- partum Stillbirths	Intra- partum Stillbirths	Not Stated	Stillbirth Rate	Early Neonatal Deaths	Early Neonatal Mortality Rate	Perinatal Mortality Rate	Adjusted PMR
Less than 22	1	0.0	6	0	1	875.0	1	1,000.0	1,000.0	1,000.0
22-27	212	0.3	61	7	4	253.5	53	250.0	440.1	417.6
28-31	515	0.7	63	0	2	112.1	16	31.1	139.7	104.1
32-36	3,422	4.8	81	4	6	25.9	41	12.0	37.6	20.0
37-41	64,951	90.8	128	16	11	2.4	21	0.3	2.7	2.0
42 and Over	2,459	3.4	3	0	0	1.2	0	0.0	1.2	0.8
Not Stated	10	0.0	0	0	0	0.0	0	0.0	0.0	0.0
Total	71,570	100.0	342	27	24	5.5	132	1.8	7.3	5.3

# TABLE REFERENCE GUIDE TO PREVIOUS PERINATAL STATISTICS REPORT 2006

#### **SECTION TWO: SELECTED TRENDS**

2007 Report	2006 Report	Title
2.1	2.2	Trends in Selected Statistics
2.2	2.1	Live Births, Stillbirths, Early Neonatal Deaths, and Perinatal Deaths, Ireland: 1999-2007
2.3	2.1	Singleton and Multiple Births: 2003-2007
2.4	2.1	Live Births, Stillbirths and Early Neonatal Deaths, Numbers and Rates, for Singleton and Multiple Births: 2003-2007

#### **SECTION THREE: GENERAL CHARACTERISTICS**

2007 Report	2006 Report	Title
3.1a	3.1	Age of Mother, Singleton Births
3.1b	3.16	Age of Mother, Multiple Births
3.2a	3.3	Mother's Marital Status, Singleton Births
3.2b	3.18	Mother's Marital Status, Multiple Births
3.3a	3.5	Mother's Occupation, Singleton Births
3.3b	3.20	Mother's Occupation, Multiple Births
3.4a	3.4	Father's Occupation, Singleton Births
3.4b	3.19	Father's Occupation, Multiple Births
3.5a	3.13	Mother's Nationality, Singleton Births
3.5b	3.28	Mother's Nationality, Multiple Births
3.6a	3.14	Father's Nationality, Singleton Births
3.6b	3.29	Father's Nationality, Multiple Births
3.7a	3.2	Maternal Parity, Singleton Births
3.7b	3.17	Maternal Parity, Multiple Births
3.8a	3.6	Previous Stillbirths, Singleton Births
3.8b	3.21	Previous Stillbirths, Multiple Births
3.9a	3.11	Interval in Years Since Last Birth, Singleton Births
3.9b	3.26	Interval in Years Since Last Birth, Multiple Births
3.10a	3.7	Birthweight, Singleton Births
3.10b	3.22	Birthweight, Multiple Births
3.11a	3.10	Gestational Age at Delivery, Singleton Births
3.11b	3.25	Gestational Age at Delivery, Multiple Births
3.12a	3.12	Month of Birth, Singleton Births
3.12b	3.27	Month of Birth, Multiple Births

## **SECTION FOUR: PERINATAL CARE**


2007 Report	2006 Report	Title
4.1a	4.1	Type of Antenatal Care, Singleton Births
4.1b	4.10	Type of Antenatal Care, Multiple Births
4.2a	l1	Time of First Visit to Doctor or Hospital During Pregnancy, Singleton Births
4.2b	12	Time of First Visit to Doctor or Hospital During Pregnancy, Multiple Births
4.3a	4.2	Mother's Antenatal Length of Stay, Singleton Births
4.3b	4.11	Mother's Antenatal Length of Stay, Multiple Births
4.4a	4.3	Mother's Postnatal Length of Stay, Singleton Births
4.4b	4.12	Mother's Postnatal Length of Stay, Multiple Births
4.5a	4.5	Advance Hospital Bookings, Singleton Births
4.5b	4.14	Advance Hospital Bookings, Multiple Births
4.6a	4.8	Rubella: Immune Status of Mother, Singleton Births
4.6b	4.16	Rubella: Immune Status of Mother, Multiple Births
4.7a	4.4	Method of Delivery, Singleton Births
4.7b	4.13	Method of Delivery, Multiple Births
4.8	N/A	Method of Delivery by Mother's Age, Total Births
4.9	4.6	Size of Maternity Unit (number of Live Births and Stillbirths Annually)
4.10a	4.7	Inter-Hospital Transfer of Infants, Singleton Births
4.10b	4.15	Inter-Hospital Transfer of Infants, Multiple Births
4.11a	5.5	Percentage Distribution of Live Births by Infant's Length of Stay, Singleton Births
4.11b	5.11	Percentage Distribution of Live Births by Infant's Length of Stay, Multiple Births
4.12a	4.9	Age of Mother by Infant's Type of Feeding, Singleton Births
4.12b	4.17	Age of Mother by Infant's Type of Feeding, Multiple Births
4.13	N/A	Mother's Occupation by Infant's Type of Feeding, Total Live Births

### **SECTION FIVE: PERINATAL OUTCOMES**

2007 Report	2006 Report	Title				
5.1a	5.1	Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal Deaths,				
		and Mortality Rates by Birthweight, Singleton Births				
5.1b	5.7	Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal Deaths, and Mortality Rates by Birthweight, Multiple Births				
5.2a	5.3	Perinatal Mortality: Numbers and Rates, by Birthweight and Cause of Death, Singleton Births				
5.2b	5.9	Perinatal Mortality: Numbers and Rates, by Birthweight and Cause of Death, Multiple Births				
5.3a	N/A	Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal Deaths, and Mortality Rates by Gestational Age, Singleton Births				
5.3b	N/A	Live Births, Antepartum and Intrapartum Stillbirths, Early Neonatal Deaths, and Mortality Rates by Gestational Age, Multiple Births				
5.4a	5.2	Stillbirths, Early Neonatal Deaths, and Mortality Rates by Cause of Death, Singleton Births				
5.4b	5.8	Stillbirths, Early Neonatal Deaths, and Mortality Rates by Cause of Death, Multiple Births				
5.5a	5.4	Age at Death, Early Neonatal Deaths, Singleton Births				
5.5b	5.10	Age at Death, Early Neonatal Deaths, Multiple Births				
5.6a	5.6	Stillbirths and Early Neonatal Deaths undergoing Port-Mortem Examinations, Singleton Births				
5.6b	5.12	Stillbirths and Early Neonatal Deaths undergoing Port-Mortem Examinations, Multiple Births				

#### **APPENDIX G**

2007 Report	2006 Report	Title
G1 – G8	H1 – H8	Selected Statistics and Cross Tabulations on Domiciliary Births



# Health Research and Information Division The Economic and Social Research Institute

Whitaker Square Sir John Rogerson's Quay Dublin 2 Ireland

Telephone + 353 1 8632000 Fax + 353 1 8632100

www.esri.ie